

INTUITIONISTIC LOGIC WITH TWO GALOIS CONNECTIONS COMBINED WITH FISCHER SERVI AXIOMS

WOJCIECH DZIK, JOUNI JÄRVINEN, AND MICHIO KONDO

ABSTRACT. Earlier, the authors introduced the logic IntGC, which is an extension of intuitionistic propositional logic by two rules of inference mimicking the performance of Galois connections (Logic J. of the IGPL, 18:837-858, 2010). In this paper, the extensions Int2GC and Int2GC+FS of IntGC are studied. Int2GC can be seen as a fusion of two IntGC logics, and Int2GC+FS is obtained from Int2GC by adding instances of duality-like connections $\Diamond(A \rightarrow B) \rightarrow (\Box A \rightarrow \Diamond B)$ and $(\Diamond A \rightarrow \Box B) \rightarrow \Box(A \rightarrow B)$, introduced by G. Fischer Servi (Rend. Sem. Mat. Univers. Politecn. Torino, 42:179-194, 1984), for interlinking the two Galois connections of Int2GC. Both Kripke-style and algebraic semantics are presented for Int2GC and Int2GC+FS, and the logics are proved to be complete with respect to both of these semantics. We show that rough lattice-valued fuzzy sets defined on complete Heyting algebras are proper algebraic models for Int2GC+FS. We also prove that Int2GC+FS is equivalent to the intuitionistic tense logic IK_t , and an axiomatisation of IK_t with the number of axioms reduced to the half of the number of axioms given by W. B. Ewald (J. Symb. Log, 51:166–179, 1986) is presented.

Key words and phrases: intuitionistic logic, Galois connections, information logic, rough lattice-valued fuzzy sets on complete Heyting algebras, Kripke semantics, intuitionistic tense logic, completeness theorems.

1. INTRODUCTION AND MOTIVATION

In [15], Information Logic of Galois Connections (ILGC) was introduced as classical propositional logic with a pair of unary connectives $\blacktriangle$ and ∇ mimicking a Galois connection. Motivation for ILGC originates in *rough set theory* [16], where it is assumed that our knowledge about objects of a universe of discourse U is expressed by an *information relation* R . An information relation may reflect similarity or difference between objects. For instance, R can be defined on the set of all human beings in such a way that two persons are R -related if they are of the same gender and the difference of their ages is less than a year. Originally, Pawlak assumed information relations to be equivalences (reflexive, symmetric, and transitive binary relations), so called *indiscernibility relations*, but in the literature can be found numerous studies considering information relations of different type; see [4], for example.

In terms of an information relation R , we may define the *upper approximation* of a set $X \subseteq U$ as

$$X^\blacktriangle = \{x \in U \mid (\exists y \in U) xRy \ \& \ y \in X\},$$

and the *lower approximation* of X is

$$X^\nabla = \{x \in U \mid (\forall y \in U) xRy \Rightarrow y \in X\}.$$

For instance, if R is the information relation considered above, then $x \in X^\nabla$ if all the persons that are coarsely of the same age and are of the same gender as x belong to X , and $x \in X^\blacktriangle$ if there exists at least one such person. Therefore, ∇ may

be interpreted to represent certainty and $\blacktriangle$ possibility with respect to knowledge expressed by the relation R .

We may also define another pair of mappings $\wp(U) \rightarrow \wp(U)$ by reversing the relation R . For any set $X \subseteq U$, let us define

$$X^\Delta = \{x \in U \mid (\exists y \in U) yRx \text{ \& } y \in X\}$$

and

$$X^\nabla = \{x \in U \mid (\forall y \in U) yRx \Rightarrow y \in X\}.$$

It is well-known that for any binary relation, the pairs $(\blacktriangle, \nabla)$ and $(\Delta, \blacktriangledown)$ are order-preserving Galois connections $\wp(U) \rightarrow \wp(U)$.

The logic **ILGC** was defined by adding to classical propositional logic two rules of inference:

$$\begin{array}{c} \text{(GC } \nabla \blacktriangle) \quad \frac{A \rightarrow \nabla B}{\blacktriangle A \rightarrow B} \qquad \qquad \text{(GC } \blacktriangle \nabla) \quad \frac{\blacktriangle A \rightarrow B}{A \rightarrow \nabla B} \end{array}$$

Another pair of connectives is introduced by De Morgan-type assertions:

$$(\star) \qquad \Delta A = \neg \nabla \neg A \quad \text{and} \quad \blacktriangledown A = \neg \blacktriangle \neg A,$$

For Δ and $\blacktriangledown$ the following rules are admissible in **ILGC**:

$$\begin{array}{c} \text{(GC } \blacktriangledown \Delta) \quad \frac{A \rightarrow \blacktriangledown B}{\Delta A \rightarrow B} \qquad \qquad \text{(GC } \Delta \blacktriangledown) \quad \frac{\Delta A \rightarrow B}{A \rightarrow \blacktriangledown B} \end{array}$$

This means that in **ILGC**, we get another Galois connection $(\Delta, \blacktriangledown)$ “for free”.

In [7], we introduced an intuitionistic propositional logic with a Galois connection (**IntGC**) and studied its main properties. In addition to the intuitionistic logic axioms and inference rule of Modus Ponens, **IntGC** contains rules (GC $\nabla \blacktriangle$) and (GC $\blacktriangle \nabla$). Since the base logic is changed from classical to intuitionistic, the classical-type assertions $(\star)$ can not be used to introduce another Galois connection. More precisely, if we define the operators $\Delta, \blacktriangledown$ from $\nabla, \blacktriangle$ in terms of intuitionistic negation, the pair $(\Delta, \blacktriangledown)$ does not form a Galois connection; see Lemma 3.3 in [7]. Therefore, to define an intuitionistic logic of two Galois connections, the other Galois connection must be declared by adding rules (GC $\blacktriangledown \Delta$) and (GC $\Delta \blacktriangledown$).

In Section 2, we define two intuitionistic logics with two Galois connections. The first one, called **Int2GC**, is obtained by extending intuitionistic propositional logic with the connectives $\blacktriangle, \blacktriangledown, \Delta, \nabla$ and by rules (GC $\nabla \blacktriangle$), (GC $\blacktriangle \nabla$), (GC $\blacktriangledown \Delta$), (GC $\Delta \blacktriangledown$). In **Int2GC**, the two Galois connections $(\blacktriangle, \nabla)$ and $(\Delta, \blacktriangledown)$ are not connected with each other, and this means that **Int2GC** is simply the fusion of two **IntGC** logics, the first one having the operators $\blacktriangle$ and ∇ , and the second has Δ and $\blacktriangledown$. The logic **Int2GC+FS** is obtained by extending **Int2GC** with instances of the axioms $\Diamond(A \rightarrow B) \rightarrow (\Box A \rightarrow \Diamond B)$ and $(\Diamond A \rightarrow \Box B) \rightarrow \Box(A \rightarrow B)$ introduced by Fischer Servi [10]. This means that **Int2GC+FS** has duality-like connections $\blacktriangle(A \rightarrow B) \rightarrow (\blacktriangledown A \rightarrow \blacktriangle B)$, $\Delta(A \rightarrow B) \rightarrow (\nabla A \rightarrow \Delta B)$, $(\blacktriangle A \rightarrow \blacktriangledown B) \rightarrow \blacktriangledown(A \rightarrow B)$, and $(\Delta A \rightarrow \nabla B) \rightarrow \nabla(A \rightarrow B)$. These axioms defining **Int2GC+FS** are referred to as (FS1), (FS2), (FS3), and (FS4), respectively. We show that in **Int2GC**, axioms (FS1) and (FS4) are equivalent, and the same holds with (FS2) and (FS3). This implies that we have several equivalent combinations of axioms to define **Int2GC+FS**.

Section 3 is devoted to **H2GC**- and **H2GC+FS**-algebras that are used for defining algebraic semantics for **Int2GC** and **Int2GC+FS**, respectively. **H2GC**-algebras are Heyting algebras equipped with two order-preserving Galois connections $(\blacktriangleright, \blacktriangleleft)$ and $(\blacktriangleright, \blacktriangleleft)$, and **H2GC+FS**-algebras are **H2GC**-algebras such that the operations $\blacktriangleright$ and $\blacktriangleleft$ are connected by an identity corresponding to axiom (FS1), and $\blacktriangleright$ and $\blacktriangleleft$ are connected by an equation that corresponds (FS2). In [6], J. M. Dunn studied distributive lattices with two operators $\Box$ and $\Diamond$. He introduced conditions $(D_\wedge) \Diamond x \wedge \Box y \leq \Diamond(x \wedge y)$ and $(D_\vee) \Box(x \vee y) \leq \Box x \vee \Diamond y$ for the interactions

between $\Box$ and $\Diamond$. We show that H2GC+FS-algebras can be defined also as H2GC-algebras satisfying the identities corresponding $(D_\wedge)$, that is, H2GC+FS-algebras are H2GC-algebras satisfying $a^\blacktriangleright \wedge b^\blacktriangleleft \rightarrow (a \wedge b)^\blacktriangleright = 1$ and $a^\blacktriangleright \wedge b^\blacktriangleleft \rightarrow (a \wedge b)^\blacktriangleright = 1$. In Section 3.2, we consider rough fuzzy sets defined on complete Heyting algebras, and show how in this setting H2GC+FS-algebras arise naturally. Algebras of rough fuzzy sets satisfy $(D_\wedge)$ when $\Diamond$ and $\Box$ are interpreted by $\blacktriangleright$ and $\blacktriangleleft$ (or $\triangleright$ and $\triangleleft$), but condition $(D_\vee)$ is not satisfied. So, rough fuzzy sets are proper algebraic models for Int2GC+FS. In Section 3.3, we introduce algebraic semantics for Int2GC and Int2GC+FS with respect to H2GC- and H2GC+FS-algebras, respectively, and present algebraic completeness theorems.

In Section 4, Kripke-semantics for Int2GC and Int2GC+FS are considered. We begin with recalling Kripke-frames and completeness for IntGC from [7] in Section 4.1. In addition, we introduce Kripke-frames and semantics for Int2GC and Int2GC+FS, and soundness of both Int2GC and Int2GC+FS is proved. Canonical frames of H2GC-algebras are introduced and Kripke-completeness is proved. Section 4.1 ends by an example in which particular Kripke-frames for Int2GC+FS are defined in terms of preference relations. In Section 4.2, we define canonical frames of H2GC+FS-algebras, and Kripke-completeness of Int2GC+FS is proved by applying canonical frames and algebraic completeness result of Int2GC+FS.

It is proved in [15] that ILGC is equivalent, with respect to provability, to the minimal (classical) tense logic K_t , that is, ILGC can be viewed as a simple formulation of K_t . In Section 5, we prove that intuitionistic tense logic IK_t , introduced by Ewald [9], is equivalent syntactically to Int2GC+FS when $\blacktriangle$, $\blacktriangledown$, Δ , ∇ are identified with tense operators F , G , P , H , respectively. In other words, in IK_t and Int2GC+FS exactly the same formulas can be proved. This then means that Int2GC+FS can be seen as an alternative formulation of IK_t . In addition, we give an axiomatisation of IK_t with the number of axioms reduced to half of the number of axioms of IK_t (with the same rules) given by Ewald [9], and we present another definition of Int2GC using only axioms of Ewald and rules admissible in IK_t .

2. INTUITIONISTIC LOGICS WITH GALOIS CONNECTIONS AND FISCHER SERVI AXIOMS

In this section we introduce two modal logics Int2GC and Int2GC+FS based on *intuitionistic propositional logic* [3, 17]. We begin with recalling the intuitionistic propositional logic with a Galois connection (IntGC) defined by the authors in [7]. The language of IntGC is constructed from an enumerable infinite set of propositional variables Var , the connectives $\neg$, $\vee$, $\wedge$, $\rightarrow$, and the unary operators $\blacktriangle$ and ∇ . The constant *true* is defined by setting $\top := p \rightarrow p$ for some fixed propositional variable $p \in \text{Var}$, and the constant *false* is defined by $\perp := \neg\top$. We also set $A \leftrightarrow B := (A \rightarrow B) \wedge (B \rightarrow A)$. The logic IntGC is the smallest logic that contains intuitionistic propositional logic, and is closed under the rules of *substitution*, *modus ponens*, and rules $(GC \nabla \blacktriangle)$ and $(GC \blacktriangle \nabla)$. The following rules are admissible in IntGC:

$$\begin{aligned} (RN\nabla) \quad & \frac{A}{\nabla A} \\ (RM\nabla) \quad & \frac{A \rightarrow B}{\nabla A \rightarrow \nabla B} \qquad (RM\blacktriangle) \quad \frac{A \rightarrow B}{\blacktriangle A \rightarrow \blacktriangle B} \end{aligned}$$

In addition, the following formulas are provable:

- (GC1) $A \rightarrow \nabla \blacktriangle A$ and $\blacktriangle \nabla A \rightarrow A$;
- (GC2) $\blacktriangle A \leftrightarrow \blacktriangle \nabla \blacktriangle A$ and $\nabla A \leftrightarrow \nabla \blacktriangle \nabla A$;
- (GC3) $\nabla \top$ and $\neg \blacktriangle \perp$;

(GC4) $\nabla(A \wedge B) \leftrightarrow \nabla A \wedge \nabla B$ and $\blacktriangle(A \vee B) \leftrightarrow \blacktriangle A \vee \blacktriangle B$;
 (GC5) $\nabla(A \rightarrow B) \rightarrow (\nabla A \rightarrow \nabla B)$.

The language of the logic **Int2GC** is the one of **IntGC** extended by two unary connectives Δ and ∇ , and the logic **Int2GC** is the smallest logic extending **IntGC** by rules (GC $\nabla\Delta$) and (GC $\Delta\nabla$). Obviously, in **Int2GC** also the rules:

$$\begin{array}{c} \text{(RN}\nabla\text{)} \frac{A}{\nabla A} \\[10pt] \text{(RM}\nabla\text{)} \frac{A \rightarrow B}{\nabla A \rightarrow \nabla B} \qquad \qquad \text{(RM}\Delta\text{)} \frac{A \rightarrow B}{\Delta A \rightarrow \Delta B} \end{array}$$

are admissible, and the following formulas are provable:

(GC1)* $A \rightarrow \nabla\Delta A$ and $\Delta\nabla A \rightarrow A$;
 (GC2)* $\Delta A \leftrightarrow \Delta\nabla\Delta A$ and $\nabla A \leftrightarrow \nabla\Delta\nabla A$;
 (GC3)* $\nabla\top$ and $\neg\Delta\perp$;
 (GC4)* $\nabla(A \wedge B) \leftrightarrow \nabla A \wedge \nabla B$ and $\Delta(A \vee B) \leftrightarrow \Delta A \vee \Delta B$;
 (GC5)* $\nabla(A \rightarrow B) \rightarrow (\nabla A \rightarrow \nabla B)$.

Intuitionistic modal logic IK was introduced by G. Fischer Servi in [10]. The logic **IK** is obtained by adding two modal connectives $\Diamond$ and $\Box$ to intuitionistic logic satisfying the following axioms:

(IK1) $\Diamond(A \vee B) \rightarrow \Diamond A \vee \Diamond B$
 (IK2) $\Box A \wedge \Box B \rightarrow \Box(A \wedge B)$
 (IK3) $\neg\Diamond\perp$
 (IK4) $\Diamond(A \rightarrow B) \rightarrow (\Box A \rightarrow \Diamond B)$
 (IK5) $(\Diamond A \rightarrow \Box B) \rightarrow \Box(A \rightarrow B)$

In addition, the *monotonicity rules* for both $\Diamond$ and $\Box$ are admissible, that is:

$$\begin{array}{c} \text{(RM}\Diamond\text{)} \frac{A \rightarrow B}{\Diamond A \rightarrow \Diamond B} \qquad \qquad \text{(RM}\Box\text{)} \frac{A \rightarrow B}{\Box A \rightarrow \Box B} \end{array}$$

In this work, we call axioms (IK4) and (IK5) the *Fischer Servi axioms*, and they have a special role in interlinking the two Galois connections of **Int2GC**. From (IK4) and (IK5) we can form the following four axioms by replacing $\Box$ and $\Diamond$ by ∇ and $\blacktriangle$, and by ∇ and Δ , respectively:

(FS1) $\blacktriangle(A \rightarrow B) \rightarrow (\nabla A \rightarrow \blacktriangle B)$
 (FS2) $\Delta(A \rightarrow B) \rightarrow (\nabla A \rightarrow \Delta B)$
 (FS3) $(\blacktriangle A \rightarrow \nabla B) \rightarrow \nabla(A \rightarrow B)$
 (FS4) $(\Delta A \rightarrow \nabla B) \rightarrow \nabla(A \rightarrow B)$

Proposition 2.1. *In Int2GC, the following assertions hold:*

- (a) *Axioms (FS1) and (FS4) are equivalent.*
- (b) *Axioms (FS2) and (FS3) are equivalent.*

Proof. We prove only assertion (a), because (b) can be proved analogously. Here $\vdash A$ denotes that A is provable in **Int2GC**.

(FS1) $\Rightarrow$ (FS4): Let us set $X := A$, $Y := \nabla\Delta A$ and $Z := \blacktriangle\nabla B$ in the provable formula $(X \rightarrow Y) \rightarrow ((Y \rightarrow Z) \rightarrow (X \rightarrow Z))$. We get $\vdash (\nabla\Delta A \rightarrow \blacktriangle\nabla B) \rightarrow (A \rightarrow \blacktriangle\nabla B)$ by using also $\vdash A \rightarrow \nabla\Delta A$. This is equivalent to $\vdash A \wedge (\nabla\Delta A \rightarrow \blacktriangle\nabla B) \rightarrow \blacktriangle\nabla B$. Because $\vdash \blacktriangle\nabla B \rightarrow B$, this means $\vdash A \wedge (\nabla\Delta A \rightarrow \blacktriangle\nabla B) \rightarrow B$ and $\vdash (\nabla\Delta A \rightarrow \blacktriangle\nabla B) \rightarrow (A \rightarrow B)$. If we set $A := \Delta A$ and $B := \nabla B$ in (FS1), we obtain $\vdash \blacktriangle(\Delta A \rightarrow \nabla B) \rightarrow (\nabla\Delta A \rightarrow \blacktriangle\nabla B)$, and so $\vdash \blacktriangle(\Delta A \rightarrow \nabla B) \rightarrow (A \rightarrow B)$. This implies $\vdash (\Delta A \rightarrow \nabla B) \rightarrow \nabla(A \rightarrow B)$ by (GC $\blacktriangle\nabla$).

(FS4) $\Rightarrow$ (FS1): We set $X := \Delta\nabla A$, $Y := A$ and $Z := B$ in $(X \rightarrow Y) \rightarrow ((Y \rightarrow Z) \rightarrow (X \rightarrow Z))$. This gives $\vdash (\Delta\nabla A \rightarrow A) \rightarrow ((A \rightarrow B) \rightarrow (\Delta\nabla A \rightarrow B))$, and $\vdash (A \rightarrow$

$B) \rightarrow (\Delta \nabla A \rightarrow \nabla \blacktriangle B)$, since $\vdash \Delta \nabla A \rightarrow A$ and $\vdash B \rightarrow \nabla \blacktriangle B$. By monotonicity, $\vdash \blacktriangle(A \rightarrow B) \rightarrow \blacktriangle(\Delta \nabla A \rightarrow \nabla \blacktriangle B)$. By setting $A := \nabla A$ and $B := \blacktriangle B$ in (FS4), we have $\vdash (\Delta \nabla A \rightarrow \nabla \blacktriangle B) \rightarrow \nabla(\nabla A \rightarrow \blacktriangle B)$ and $\vdash \blacktriangle(\Delta \nabla A \rightarrow \nabla \blacktriangle B) \rightarrow (\nabla A \rightarrow \blacktriangle B)$ by (GC $\nabla \blacktriangle$). Therefore, we obtain $\vdash \blacktriangle(A \rightarrow B) \rightarrow (\nabla A \rightarrow \blacktriangle B)$. $\square$

The logic $\text{Int2GC}+\text{FS}$ is defined as the extension of Int2GC that satisfies also the Fischer Servi axioms (FS1)–(FS4). By Proposition 2.1 it is clear that we have several equivalent axiomatisations of $\text{Int2GC}+\text{FS}$, that is:

$$\text{Int2GC}+\text{FS} = \text{Int2GC} + \{(\text{FS1}) \text{ or } (\text{FS4})\} + \{(\text{FS2}) \text{ or } (\text{FS3})\}.$$

The logic $\text{Int2GC}+\text{FS}$ satisfies the counterparts of axioms (IK1)–(IK5) of IK, so $\text{Int2GC}+\text{FS}$ can be regarded as a intuitionistic bi-modal logic, and the pairs $\blacktriangle$, ∇ and Δ , ∇ are intuitionistic modal connectives in the sense of Fischer Servi.

3. ALGEBRAIC SEMANTICS AND COMPLETENESS

3.1. H2GC-algebras and H2GC+FS-algebras. In [7], we introduced HGC-algebras as counterparts of the logic IntGC , and we showed that IntGC is complete with respect to HGC-algebras. In this section, we define H2GC- and H2GC+FS-algebras and give completeness theorems for Int2GC and $\text{Int2GC}+\text{FS}$ in terms of these algebras.

Let $\varphi: P \rightarrow Q$ and $\psi: Q \rightarrow P$ be maps between ordered sets P and Q . The pair (φ, ψ) is a *Galois connection* between P and Q , if for all $p \in P$ and $q \in Q$,

$$\varphi(p) \leq q \iff p \leq \psi(q).$$

For a Galois connection (φ, ψ) , φ preserves all existing joins and ψ preserves all existing meets. If P and Q are bounded, then $\varphi(0) = 0$ and $\psi(1) = 1$. In addition, a pair (φ, ψ) forms a Galois connection if and only if the following conditions hold:

- (i) $p \leq \psi(\varphi(p))$ for all $p \in P$ and $\varphi(\psi(q)) \leq q$ for all $q \in Q$;
- (ii) the maps φ and ψ are order-preserving.

In particular, if φ and ψ are maps on a lattice L , then the pair (φ, ψ) is a Galois connection on L if and only if the following identities are satisfied for all $a, b \in L$:

- (gc1) $\varphi(a \vee b) = \varphi(a) \vee \varphi(b)$ and $\psi(a \wedge b) = \psi(a) \wedge \psi(b)$
- (gc2) $a = a \wedge \psi(\varphi(a))$ and $a = a \vee \varphi(\psi(a))$

More properties of Galois connections can be found in [8], for instance.

A *Heyting algebra* H is a lattice with 0 such that for all $a, b \in H$, there is a greatest element x of H with $a \wedge x \leq b$. This element is the *relative pseudocomplement* of a with respect to b , and is denoted $a \rightarrow b$. Note that Heyting algebras are always distributive pseudocomplemented lattices such that the pseudocomplement $\neg a$ of a is $a \rightarrow 0$. Because $\neg 0$ is the greatest element, Heyting algebras are bounded. Thus, a Heyting algebra H can be considered as an algebraic structure $\mathbb{H} = (H, \vee, \wedge, \rightarrow, 0)$, which can be equationally defined as follows (see e.g. [1]):

- (h1) A set of identities which define lattice with 0
- (h2) $x \wedge (x \rightarrow y) = x \wedge y$
- (h3) $x \wedge (y \rightarrow z) = x \wedge ((x \wedge y) \rightarrow (x \wedge z))$
- (h4) $z \wedge (x \wedge y \rightarrow x) = x$

Note also that if $\mathbb{H}$ is a Heyting algebra, then (gc2) can be written in the form

$$(\text{gc2})^* \quad a \rightarrow \psi(\varphi(a)) = 1 \text{ and } \varphi(\psi(a)) \rightarrow a = 1.$$

An *HGC-algebra* is an algebra $(H, \vee, \wedge, \rightarrow, 0, \blacktriangleright, \blacktriangleleft)$, where $\mathbb{H} = (H, \vee, \wedge, \rightarrow, 0)$ is a Heyting algebra and $(\blacktriangleright, \blacktriangleleft)$ is a Galois connection on $\mathbb{H}$. By the above, HGC-algebras form an equational class. HGC-algebras are usually denoted by $(\mathbb{H}, \blacktriangleright, \blacktriangleleft)$. An *H2GC-algebra* $(H, \vee, \wedge, \rightarrow, 0, \blacktriangleright, \blacktriangleleft, \triangleright, \triangleleft)$ is such that $\mathbb{H} = (H, \vee, \wedge, \rightarrow, 0)$ is a Heyting algebra, and $(\mathbb{H}, \blacktriangleright, \blacktriangleleft)$ and $(\mathbb{H}, \triangleright, \triangleleft)$ are H2GC-algebras, meaning that

$(\blacktriangleright, \blacktriangleleft)$ and $(\blacktriangleright, \blacktriangleleft)$ are Galois connections on $\mathbb{H}$. Also H2GC-algebras form an equational class. We denote H2GC-algebras simply by $(\mathbb{H}, \blacktriangleright, \blacktriangleleft, \blacktriangleright, \blacktriangleleft)$.

For an H2GC-algebra $(\mathbb{H}, \blacktriangleright, \blacktriangleleft, \blacktriangleright, \blacktriangleleft)$, we introduce the following identities corresponding to the instances (FS1)–(FS4) of the Fischer Servi axioms:

- (fs1) $(a \rightarrow b) \blacktriangleright \rightarrow (a \blacktriangleleft \rightarrow b \blacktriangleright) = 1$
- (fs2) $(a \rightarrow b) \blacktriangleright \rightarrow (a \blacktriangleleft \rightarrow b \blacktriangleright) = 1$
- (fs3) $(a \blacktriangleright \rightarrow b \blacktriangleleft) \rightarrow (a \rightarrow b) \blacktriangleleft = 1$
- (fs4) $(a \blacktriangleright \rightarrow b \blacktriangleleft) \rightarrow (a \rightarrow b) \blacktriangleleft = 1$

In [6], J. M. Dunn studied minimal positive modal logic K_+ with the connectives $\wedge$, $\vee$, $\Box$, and $\Diamond$. K_+ can be described in algebraic terms as modal logic based on a distributive lattice with two operations $\Box$ and $\Diamond$, where $\Box$ distributes over $\wedge$, $\Diamond$ distributes over $\vee$, and the following two conditions hold:

- (D $_{\wedge}$) $\Diamond a \wedge \Box b \leq \Diamond(a \wedge b)$
- (D $_{\vee}$) $\Box(a \vee b) \leq \Box a \vee \Box b$

We introduce the instances of (D $_{\wedge}$) as identities defined on an H2GC-algebra:

- (d1) $a \blacktriangleright \wedge b \blacktriangleleft \rightarrow (a \wedge b) \blacktriangleright = 1$
- (d2) $a \blacktriangleright \wedge b \blacktriangleleft \rightarrow (a \wedge b) \blacktriangleright = 1$

Now we may write the following proposition.

Proposition 3.1. *Let $(\mathbb{H}, \blacktriangleright, \blacktriangleleft, \blacktriangleright, \blacktriangleleft)$ be an H2GC-algebra.*

- (a) *Identities (fs1), (d1), and (fs4) are equivalent.*
- (b) *Identities (fs2), (d2), and (fs3) are equivalent.*

Proof. (a) Let us set $a := a \rightarrow b$ and $b := a$ in (d1). We obtain $(a \rightarrow b) \blacktriangleright \wedge a \blacktriangleleft \leq (a \wedge (a \rightarrow b)) \blacktriangleright \leq b \blacktriangleright$, because $a \wedge (a \rightarrow b) \leq b$. This gives directly $(a \rightarrow b) \blacktriangleright \leq a \blacktriangleleft \rightarrow b \blacktriangleright$, that is, $(a \rightarrow b) \blacktriangleright \rightarrow (a \blacktriangleleft \rightarrow b \blacktriangleright) = 1$, and so (d1) implies (fs1). Conversely, if we set $b := a \wedge b$ in (fs1), we have $b \blacktriangleright \leq (a \rightarrow b) \blacktriangleright \leq (a \rightarrow a \wedge b) \blacktriangleright \leq a \blacktriangleleft \rightarrow (a \wedge b) \blacktriangleright$, because $a \rightarrow a \wedge b = a \rightarrow b$ and $b \leq a \rightarrow b$. This is equivalent to $a \blacktriangleleft \wedge b \blacktriangleright \leq (a \wedge b) \blacktriangleleft$, and $(a \blacktriangleleft \wedge b \blacktriangleright) \rightarrow (a \wedge b) \blacktriangleleft = 1$. Thus, also (fs1) implies (d1), and (fs1) and (d1) are equivalent.

Suppose that (fs1) holds. Then $(a \blacktriangleright \rightarrow b \blacktriangleleft) \blacktriangleright \leq a \blacktriangleright \blacktriangleleft \rightarrow b \blacktriangleleft \blacktriangleright$. Since $a \leq a \blacktriangleright \blacktriangleleft$ and $b \blacktriangleleft \blacktriangleright \leq b$, we have $(a \blacktriangleright \rightarrow b \blacktriangleleft) \blacktriangleright \leq a \rightarrow b$. This is equivalent to $a \blacktriangleright \rightarrow b \blacktriangleleft \leq (a \rightarrow b) \blacktriangleleft$ and $(a \blacktriangleright \rightarrow b \blacktriangleleft) \rightarrow (a \rightarrow b) \blacktriangleleft = 1$, that is, (fs4) is true. On the other hand, if (fs4) holds, then $a \rightarrow b \leq a \blacktriangleleft \blacktriangleright \rightarrow b \blacktriangleright \blacktriangleleft \leq (a \blacktriangleleft \rightarrow b \blacktriangleright) \blacktriangleleft$, that is, $(a \rightarrow b) \blacktriangleright \leq a \blacktriangleleft \rightarrow b \blacktriangleright$, $(a \rightarrow b) \blacktriangleright \leq (a \blacktriangleleft \rightarrow b \blacktriangleright) = 1$, and (fs1) is true. Hence, (fs1) and (fs4) are equivalent.

Case (b) can be proved analogously. $\square$

An *H2GC+FS-algebra* is an H2GC-algebra $(\mathbb{H}, \blacktriangleright, \blacktriangleleft, \blacktriangleright, \blacktriangleleft)$ satisfying (fs1) and (fs2). By Proposition 3.1, H2GC+FS-algebras have several equivalent characterisations. Clearly, H2GC+FS-algebras form an equational class.

3.2. Rough fuzzy sets on complete Heyting algebras. We consider here rough lattice-valued fuzzy sets defined on complete Heyting algebras. These are also closely connected to fuzzy Galois connections studied, for instance, in [2, 11].

A *complete Heyting algebra* is a Heyting algebra such that its underlying ordered set is a complete lattice. A complete lattice L satisfies the *join-infinite distributive law* if for any $S \subseteq L$ and $x \in L$,

$$(JID) \quad x \wedge \left(\bigvee S \right) = \bigvee \{x \wedge y \mid y \in S\}.$$

A complete lattice is a Heyting algebra if and only if it satisfies (JID) (see e.g. [13, 17]). Thus, complete Heyting algebras are the complete lattices satisfying (JID).

Fuzzy sets were generalized to L -fuzzy sets by J. A. Goguen in such a way that an L -fuzzy set φ on U is a mapping $\varphi: U \rightarrow L$, where U is any set representing

objects of some universe of discourse and L is a partially ordered set [12]. The set L^U of all maps from U to L is then the set of all L -fuzzy sets on U . The set L^U can be equipped whatever operators L has, and these induced operators obey any law valid in L which extends pointwise.

Here we assume that $\mathbb{H}$ is a complete Heyting algebra, therefore we can make H^U a complete Heyting algebra by defining

$$\left(\bigvee_{i \in I} \varphi_i\right)(a) = \bigvee_{i \in I} \varphi_i(a) \quad \text{and} \quad \left(\bigwedge_{i \in I} \varphi_i\right)(a) = \bigwedge_{i \in I} \varphi_i(a)$$

for all $\{\varphi_i\}_{i \in I} \subseteq H^U$. The least element of H^U is $\mathbf{0}: x \mapsto 0$ and the greatest element of H^U is $\mathbf{1}: x \mapsto 1$. Furthermore, H^U is relatively pseudocomplemented in such a way that for all $\varphi, \psi \in L^U$ and $a \in U$,

$$(\varphi \rightarrow \psi)(a) = \varphi(a) \rightarrow \psi(a)$$

We denote this complete Heyting algebra by $\mathbb{H}^U$. Elements of this Heyting algebra are called $\mathbb{H}$ -sets.

Dubois and Prade introduced *rough fuzzy sets* in [5]. The idea is that the objects to be approximated are fuzzy sets, and the approximations are determined by means of fuzzy relations. Here we study *rough $\mathbb{H}$ -sets*, which means that approximations of $\mathbb{H}$ -sets are determined by $\mathbb{H}$ -fuzzy relations.

Let φ be an $\mathbb{H}$ -set and let R be an $\mathbb{H}$ -fuzzy relation on U , that is, R is a mapping from $U \times U$ to H . Then, we may define the $\mathbb{H}$ -sets $\varphi^\blacktriangle$ and $\varphi^\blacktriangledown$ by setting

$$\begin{aligned} \varphi^\blacktriangle(x) &= \bigvee_{y \in U} \{R(x, y) \wedge \varphi(y)\} \\ \varphi^\blacktriangledown(x) &= \bigwedge_{y \in U} \{R(x, y) \rightarrow \varphi(y)\} \end{aligned}$$

for all $x \in U$. The $\mathbb{H}$ -sets $\varphi^\blacktriangle$ and $\varphi^\blacktriangledown$ are called the *upper* and the *lower approximations* of φ .

We can define another pair of mappings in terms of the inverse of R by setting

$$\begin{aligned} \varphi^\Delta(x) &= \bigvee_{y \in U} \{R(y, x) \wedge \varphi(y)\} \\ \varphi^\nabla(x) &= \bigwedge_{y \in U} \{R(y, x) \rightarrow \varphi(y)\} \end{aligned}$$

for all $x \in U$. It is clear that if φ is a two-valued set on U and R is a two-valued binary relation on U , then the operations $\blacktriangle$, $\blacktriangledown$, Δ , and ∇ coincide with the rough set operators defined by a binary relation.

Proposition 3.2. *For any complete Heyting algebra $\mathbb{H}$ and an $\mathbb{H}$ -fuzzy relation R on U , the algebra of rough $\mathbb{H}$ -sets $(\mathbb{H}^U, \blacktriangle, \blacktriangledown, \Delta, \nabla)$ is an $H2GC+FS$ -algebra.*

Proof. Suppose φ and ψ are $\mathbb{H}$ -sets such that $\varphi \leq \psi$. Then for all $y \in U$, $R(x, y) \wedge \varphi(y) \leq R(x, y) \wedge \psi(y)$ and this implies

$$\varphi^\blacktriangle(x) = \bigvee_{y \in U} \{R(x, y) \wedge \varphi(y)\} \leq \bigvee_{y \in U} \{R(x, y) \wedge \psi(y)\} = \psi^\blacktriangle(x).$$

Similarly, $R(y, x) \rightarrow \varphi(y) \leq R(y, x) \rightarrow \psi(y)$ for all $y \in U$. Thus,

$$\varphi^\blacktriangledown(x) = \bigwedge_{y \in U} \{R(y, x) \rightarrow \varphi(y)\} \leq \bigwedge_{y \in U} \{R(y, x) \rightarrow \psi(y)\} = \psi^\blacktriangledown(x).$$

So, $\blacktriangle$ and ∇ are order-preserving. By definition, for all $x \in U$,

$$\begin{aligned}\varphi^{\nabla\blacktriangle}(x) &= \bigvee_{y \in U} \{R(x, y) \wedge \varphi^{\nabla}(y)\} = \bigvee_{y \in U} \left\{ R(x, y) \wedge \bigwedge_{z \in U} \{R(z, y) \rightarrow \varphi(z)\} \right\} \\ &\leq \bigvee_{y \in U} \{R(x, y) \wedge (R(x, y) \rightarrow \varphi(x))\} \leq \bigvee_{y \in U} \{\varphi(x)\} = \varphi(x).\end{aligned}$$

This means that $\varphi^{\blacktriangle\nabla} \leq \varphi$. Analogously, for any $x \in U$,

$$\begin{aligned}\varphi^{\blacktriangle\nabla}(x) &= \bigwedge_{y \in U} \{R(y, x) \rightarrow \varphi^{\blacktriangle}(y)\} = \bigwedge_{y \in U} \left\{ R(y, x) \rightarrow \bigvee_{z \in U} \{R(y, z) \wedge \varphi(z)\} \right\} \\ &\geq \bigwedge_{y \in U} \{R(y, x) \rightarrow (R(y, x) \wedge \varphi(x))\} \geq \bigwedge_{y \in U} \{\varphi(x)\} = \varphi(x).\end{aligned}$$

Thus, also $\varphi \leq \varphi^{\blacktriangle\nabla}$. We have that $(\blacktriangle, \nabla)$ is a Galois connection, because (gc1) and (gc2) are satisfied. Similarly, we can show that $(\blacktriangle, \blacktriangledown)$ is a Galois connection.

Next we show that (d1) holds. For all $x, y \in U$, we have

$$\begin{aligned}R(x, y) \wedge \varphi(y) \wedge \psi(x)^{\blacktriangledown} &= R(x, y) \wedge \varphi(y) \wedge \bigwedge_{z \in U} \{R(x, z) \rightarrow \psi(z)\} \\ &\leq R(x, y) \wedge \varphi(y) \wedge (R(x, y) \rightarrow \psi(y)) \\ &= (R(x, y) \wedge (R(x, y) \rightarrow \psi(y))) \wedge \varphi(y) \\ &= R(x, y) \wedge \psi(y) \wedge \varphi(y) \\ &= R(x, y) \wedge (\varphi \wedge \psi)(y) \\ &\leq \bigvee_{z \in U} \{R(x, z) \wedge (\varphi \wedge \psi)(z)\} \\ &= (\varphi \wedge \psi)^{\blacktriangle}(x).\end{aligned}$$

Hence, for all $y \in U$,

$$R(x, y) \wedge \varphi(y) \wedge \psi(x)^{\blacktriangledown} \leq (\varphi \wedge \psi)^{\blacktriangle}(x).$$

Because complete Heyting algebras satisfy the join-infinite distributive law, we have that for all $x \in U$,

$$\begin{aligned}(\varphi^{\blacktriangle} \wedge \psi^{\blacktriangledown})(x) &= \varphi^{\blacktriangle}(x) \wedge \psi^{\blacktriangledown}(x) = \bigvee_{y \in U} \{R(x, y) \wedge \varphi(y)\} \wedge \psi^{\blacktriangledown}(x) \\ &= \bigvee_{y \in U} \{R(x, y) \wedge \varphi(y) \wedge \psi^{\blacktriangledown}(x)\} \leq (\varphi \wedge \psi)^{\blacktriangle}(x).\end{aligned}$$

Thus, $\varphi^{\blacktriangle} \wedge \psi^{\blacktriangledown} \leq (\varphi \wedge \psi)^{\blacktriangle}$. Assertion (d2) can be proved similarly. $\square$

Example 3.3. The instances

$$(a \vee b)^{\blacktriangleleft} \leq a^{\blacktriangleleft} \vee b^{\blacktriangleright} \quad \text{and} \quad (a \vee b)^{\blacktriangleleft} \leq a^{\blacktriangleleft} \vee b^{\blacktriangleright}$$

of Dunn's axiom $(D_{\vee})$ are false in some H2GC+FS-algebras of rough $\mathbb{H}$ -sets.

Namely, let $U = \{x, y\}$ and consider the finite (and hence complete) Heyting algebra $\mathbf{2}^2 \oplus 1$, that is, $\mathbb{H} = \{0, a, b, c, 1\}$ the Heyting algebra with the order $0 < a, b < c < 1$, where a and b are incomparable. Note that $\neg a = b$ and $\neg b = a$.

We define two $\mathbb{H}$ -sets φ, ψ on U by setting $\varphi(u) = 0$ and $\psi(u) = 1$ for all $u \in U$. An $\mathbb{H}$ -fuzzy relation $R: U \times U \rightarrow \mathbb{H}$ is defined by $R(x, x) = R(y, y) = a$ and

$R(x, y) = R(y, x) = b$. Then,

$$\begin{aligned} (\varphi \vee \psi)^\nabla(x) &= \bigwedge_{u \in U} (R(x, u) \rightarrow (\varphi \vee \psi)(u)) = \bigwedge_{u \in U} (R(x, u) \rightarrow (\varphi(u) \vee \psi(u))) \\ &= \bigwedge_{u \in U} (R(x, u) \rightarrow 1) = (a \rightarrow 1) \wedge (b \rightarrow 1) = 1 \wedge 1 = 1, \end{aligned}$$

but

$$\begin{aligned} \varphi^\nabla(x) \vee \psi^\blacktriangle(x) &= \bigwedge_{u \in U} (R(x, u) \rightarrow \varphi(u)) \vee \bigvee_{u \in U} (R(x, u) \wedge \psi(u)) \\ &= \bigwedge_{u \in U} (R(x, u) \rightarrow 0) \vee \bigvee_{u \in U} (R(x, u) \wedge 1) \\ &= \bigwedge_{u \in U} \neg R(x, u) \vee \bigvee_{u \in U} R(x, u) \\ &= (\neg R(x, x) \wedge \neg R(x, y)) \vee (R(x, x) \vee R(x, y)) \\ &= (\neg a \wedge \neg b) \vee (a \vee b) = 0 \vee c = c. \end{aligned}$$

Hence condition $(\varphi \vee \psi)^\blacktriangleleft \leq \varphi^\blacktriangleleft \vee \psi^\blacktriangleright$ is not satisfied, because $1 \not\leq c$. Similarly, we can show that

$$(\varphi \vee \psi)^\nabla(y) = 1 \quad \text{and} \quad \varphi^\nabla(y) \vee \psi^\blacktriangle(y) = 0 \vee c = c,$$

that is, $(\varphi \vee \psi)^\blacktriangleleft \leq \varphi^\blacktriangleleft \vee \psi^\blacktriangleright$ is not satisfied.

Hence, we may conclude this subsection by stating that the rough lattice-valued fuzzy sets defined on complete Heyting algebras are algebraic models for $\text{Int2GC}+\text{FS}$.

3.3. Algebraic Semantics and Completeness. As we already noted, IntGC is complete with respect to H2GC -algebras. Here we show completeness of Int2GC and $\text{IntGC}+\text{FS}$ with respect to H2GC - and $\text{H2GC}+\text{FS}$ -algebras.

Let $(\mathbb{H}, \blacktriangleright, \blacktriangleleft, \triangleright, \triangleleft)$ be an H2GC -algebra, where $\mathbb{H} = (H, \vee, \wedge, \rightarrow, 0)$. A *valuation* is a function $v: \text{Var} \rightarrow H$ assigning to each propositional variable p an element $v(p)$ of H . Let Φ denote the set of well-formed Int2GC -formulas. Clearly, Φ is the set of well-formed $\text{Int2GC}+\text{FS}$ -formulas as well, because these logics have the same language. The valuation v can be extended to the set Φ inductively¹:

$$\begin{aligned} v(\neg A) &= v(A) \rightarrow 0 & v(A \rightarrow B) &= v(A) \rightarrow v(B) \\ v(A \wedge B) &= v(A) \wedge v(B) & v(A \vee B) &= v(A) \vee v(B) \\ v(\blacktriangle A) &= v(A)^\blacktriangleright & v(\blacktriangledown A) &= v(A)^\blacktriangleleft \\ v(\triangle A) &= v(A)^\triangleright & v(\nabla A) &= v(A)^\triangleleft \end{aligned}$$

An Int2GC -formula A is *valid* if $v(A) = 1$ for any valuation v on any H2GC -algebra. Similarly, we may define validity of $\text{Int2GC}+\text{FS}$ -formulas over $\text{H2GC}+\text{FS}$ -algebras.

Theorem 3.4 (Soundness I).

- (a) *Provable Int2GC -formulas are valid in H2GC -algebras*
- (b) *Provable $\text{Int2GC}+\text{FS}$ -formulas are valid in $\text{H2GC}+\text{FS}$ -algebras.*

Proof. The proof concerning intuitionistic logic is standard (see [17], for instance). As we have proved in [7] for IntGC , rules $(\text{GC } \nabla \blacktriangle)$, $(\text{GC } \blacktriangle \nabla)$, $(\text{GC } \blacktriangledown \triangle)$, $(\text{GC } \triangle \blacktriangledown)$ preserve validity. Thus, (a) holds. For (b), it is clear that axioms (FS1) and (FS2) are valid, because $\text{H2GC}+\text{FS}$ -algebras satisfy identities (fs1) and (fs2). $\square$

¹Note that the idea is that the operations $\blacktriangleright, \blacktriangleleft, \triangleright, \triangleleft$ may be obtained from their logical counterparts $\blacktriangle, \blacktriangledown, \triangle, \nabla$ just by turning them 90 degrees clockwise.

To obtain completeness, we apply Lindenbaum–Tarski algebras. We denote by Φ the *algebra of Φ -formulas*, that is, the abstract algebra

$$\Phi = (\Phi, \vee, \wedge, \rightarrow, \perp, \blacktriangle, \blacktriangledown, \triangle, \nabla).$$

We define two equivalences $\equiv_1$ and $\equiv_2$ on Φ :

$$A \equiv_1 B \text{ if and only if } A \leftrightarrow B \text{ is provable in Int2GC;}$$

$$A \equiv_2 B \text{ if and only if } A \leftrightarrow B \text{ is provable in Int2GC+FS.}$$

Concerning $\vee, \wedge$, and $\rightarrow$, the next result is known from the theory of intuitionistic logic, and for $\blacktriangle, \blacktriangledown, \triangle$, and ∇ the claim follows from monotonicity.

Lemma 3.5. *The equivalences $\equiv_1$ and $\equiv_2$ are congruences on Φ .*

For any $A \in \Phi$, we denote by $[A]_1$ and $[A]_2$ the congruence class of A with respect to the congruences $\equiv_1$ and $\equiv_2$. The sets of $\equiv_1$ - and $\equiv_2$ -classes are denoted by $\Phi/\equiv_1$ and $\Phi/\equiv_2$. Next we define the *quotient algebras* of Φ with respect to $\equiv_1$ and $\equiv_2$ by introducing the following operations on $\Phi/\equiv_i$ for $i = 1, 2$:

$$\begin{aligned} [A]_i \vee_i [B]_i &= [A \vee B]_i & [A]_i \wedge_i [B]_i &= [A \wedge B]_i \\ [A]_i \rightarrow_i [B]_i &= [A \rightarrow B]_i & \mathbf{0}_i &= [\perp]_i \\ [A]_i \blacktriangleright_i &= [\blacktriangle A]_i & [A]_i \blacktriangleleft_i &= [\blacktriangledown A]_i \\ [A]_i \triangleright_i &= [\triangle A]_i & [A]_i \triangleleft_i &= [\nabla A]_i \end{aligned}$$

As we have noted, H2GC- and H2GC+FS-algebras form equational classes. By the theory of intuitionistic logic, $\Phi/\equiv_1$ satisfies the identities defining Heyting algebras. Also (gc1) and (gc2) hold by (GC1), (GC1)*, (GC4), (GC4)*. Since also identities (fs1) and (fs2) are the counterparts of axioms (FS1) and (FS2), we may write the following proposition.

Proposition 3.6.

- (a) *The algebra $(\Phi/\equiv_1, \vee_1, \wedge_1, \rightarrow_1, \mathbf{0}_1, \blacktriangleright_1, \blacktriangleleft_1, \triangleright_1, \triangleleft_1)$ is an H2GC-algebra.*
- (b) *The algebra $(\Phi/\equiv_2, \vee_2, \wedge_2, \rightarrow_2, \mathbf{0}_2, \blacktriangleright_2, \blacktriangleleft_2, \triangleright_2, \triangleleft_2)$ is an H2GC+FS-algebra.*

We define two valuations $v_1: \text{Var} \rightarrow \Phi/\equiv_1$ and $v_2: \text{Var} \rightarrow \Phi/\equiv_2$ by:

$$v_1(p) = [p]_1 \quad \text{and} \quad v_2(p) = [p]_2.$$

By a straightforward formula induction we see that $v_1(A) = [A]_1$ and $v_2(A) = [A]_2$ for all formulas $A \in \Phi$. We can now write the following results.

Lemma 3.7. *For any formula $A \in \Phi$:*

- (a) *A is provable in Int2GC if and only if $v_1(A) = \mathbf{1}$.*
- (b) *A is provable in Int2GC+FS if and only if $v_2(A) = \mathbf{1}$.*

Theorem 3.8 (Completeness I). *For any formula $A \in \Phi$:*

- (a) *A is provable in Int2GC if and only if A valid in H2GC-algebras.*
- (b) *A is provable in Int2GC+FS if and only if A valid in H2GC+FS-algebras.*

Proof. (a) Suppose that A is valid in H2GC-algebras. We have $v_1(A) = \mathbf{1}$ in $\Phi/\equiv_1$, that is, A is provable. The other direction is proved in Theorem 3.4. For (b), the proof is basically the same. $\square$

Clearly, rough $\mathbb{H}$ -sets considered in Section 3.2 provide algebraic models for Int2GC+FS. Let us introduce axioms (D1) and (D2) corresponding to equations (d1) and (d2):

$$(D1) \quad \blacktriangle A \wedge \blacktriangledown B \rightarrow \blacktriangle(A \wedge B)$$

$$(D2) \quad \triangle A \wedge \nabla B \rightarrow \triangle(A \wedge B)$$

By Proposition 3.1 and Theorem 3.8, we can write the following corollary giving additional ways to axiomatize $\text{Int2GC}+\text{FS}$.

Corollary 3.9.

$$\text{Int2GC}+\text{FS} = \text{Int2GC} + \{(\text{FS1}) \text{ or } (\text{FS4}) \text{ or } (\text{D1})\} + \{(\text{FS2}) \text{ or } (\text{FS3}) \text{ or } (\text{D2})\}$$

By Example 3.3 and Theorem 3.8, formulas $\nabla(A \vee B) \rightarrow \nabla A \vee \blacktriangle B$ and $\nabla(A \vee B) \rightarrow \nabla A \vee \triangle B$ corresponding to Dunn's condition $(D_\vee)$ are not provable in $\text{Int2GC}+\text{FS}$.

4. KRIPKE SEMANTICS AND COMPLETENESS

4.1. Kripke frames. In this section we consider Kripke frames and models for the three systems IntGC , Int2GC , and $\text{Int2GC}+\text{FS}$, where the system $\text{Int2GC}+\text{FS}$ will be shown in Section 5 to be equivalent to intuitionistic temporal logic IK_t .

IntGC-frames. A structure $\mathcal{F} = (X, \leq, R)$ is called a *Kripke frame of IntGC* (an *IntGC-frame*, in short) [7], if X is a non-empty set, $\leq$ is a preorder (reflexive and transitive binary relation) on X , and R is a relation on X such that

$$(R1) \quad (\geq \circ R \circ \geq) \subseteq R.$$

Let v be a function $v: P \rightarrow \wp(X)$ assigning to each propositional variable p a subset $v(p)$ of X with the property that $x \in v(p)$ and $x \leq y$ imply $y \in v(p)$, that is, $v(p)$ is $\leq$ -closed. Such functions are called *valuations* and the pair $\mathcal{M} = (\mathcal{F}, v)$ is called an *IntGC-model*. For any $x \in X$ and $A \in \Phi$, we define the *satisfiability relation* in $\mathcal{M}$ inductively by the following way:

$$\begin{aligned} x \models p &\iff x \in v(p), \\ x \models A \wedge B &\iff x \models A \text{ and } x \models B, \\ x \models A \vee B &\iff x \models A \text{ or } x \models B, \\ x \models A \rightarrow B &\iff \text{for all } y \geq x, y \models A \text{ implies } y \models B, \\ x \models \neg A &\iff \text{for no } y \geq x \text{ does } y \models A, \\ x \models \blacktriangle A &\iff \text{exists } y \text{ such that } x R y \text{ and } y \models A, \text{ and} \\ x \models \nabla A &\iff \text{for all } y, y R x \text{ implies } y \models A. \end{aligned}$$

Note that the satisfiability relation $\models$ is *persistent*, that is, for all formulas A , if $x \models A$ and $x \leq y$, then $y \models A$. An IntGC -formula A is *valid in a model* $\mathcal{M}$, if $x \models A$ for all $x \in X$. The formula A is *valid in a frame* $\mathcal{F}$, if A is valid in every model based on $\mathcal{F}$. A formula is *Kripke-valid* if it is valid in every frame.

We noted in [7] that IntGC is *Kripke-sound*, that is, every provable IntGC -formula is Kripke-valid. We also proved Kripke-completeness by applying canonical frames, and next we shortly recall these constructions, because a similar technique will be used later in cases of Int2GC and $\text{Int2GC}+\text{FS}$.

For an HGC-algebra $(\mathbb{H}, \blacktriangleright, \triangleleft)$, its *canonical frame* is a triple $(X_H, \subseteq, R^H)$ such that X_H is the set of the prime filters of the lattice H and the relation R^H is defined by

$$(x, y) \in R^H \iff y \subseteq [x]^{\blacktriangleright^{-1}},$$

where $[x]^{\blacktriangleright^{-1}} = \{a \in H \mid a^{\blacktriangleright} \in x\}$. The relation R^H can be described also in terms of the map $\triangleleft$ by

$$(x, y) \in R^H \iff [y]^{\triangleleft^{-1}} \subseteq x,$$

where $[y]^{\triangleleft^{-1}} = \{a \in H \mid a^{\triangleleft} \in y\}$.

For a Heyting algebra $\mathbb{H}$, we denote by $\mathcal{O}(H)$ the set of all $\leq$ -closed subsets.

Lemma 4.1. *Let $(\mathbb{H}, \blacktriangleright, \triangleleft)$ be an HGC-algebra. The pair $(\triangleleft^{-1}, \blacktriangleright^{-1})$ is a Galois connection on $(\mathcal{O}(H), \subseteq)$.*

Proof. Let $x, y \in \mathcal{O}(H)$. Suppose $[x]^{\triangleleft^{-1}} \subseteq y$. If $a \in x$, then $a \leq a^{\triangleright^{\triangleleft}}$ implies $a^{\triangleright^{\triangleleft}} \in x$, because $x \in \mathcal{O}(H)$. This means that $a^{\triangleright} \in [x]^{\triangleleft^{-1}} \subseteq y$ and so $a \in [y]^{\triangleright^{-1}}$. Therefore, $x \subseteq [y]^{\triangleright^{-1}}$. Conversely, assume that $x \subseteq [y]^{\triangleright^{-1}}$. If $a \in [x]^{\triangleleft^{-1}}$, then $a^{\triangleleft} \in x \subseteq [y]^{\triangleright^{-1}}$, that is, $a^{\triangleleft} \in y$. Because $a^{\triangleleft} \leq a$, we have $a \in y$, since $y \in \mathcal{O}(H)$. Thus, $[x]^{\triangleleft^{-1}} \subseteq y$. $\square$

Let $(\mathbb{H}, \triangleright, \triangleleft)$ be an HGC-algebra. In [7], we showed that the canonical frame $\mathcal{F}^H = (X_H, \subseteq, R^H)$ is an IntGC-frame. Let $v: \text{Var} \rightarrow H$ be a valuation on this HGC-algebra. We may now define a valuation $v^*: \text{Var} \rightarrow X_H$ for the canonical frame $\mathcal{F}^H$ by setting $x \in v^*(p)$ if and only if $v(p) \in x$ for all $p \in \text{Var}$. Obviously, for all $x, y \in X_H$ and $p \in \text{Var}$, $x \in v^*(p)$ and $x \subseteq y$ imply $y \in v^*(p)$, so v^* is really a valuation. In the *canonical model* $(\mathcal{F}^H, v^*)$, we have $x \models p$ if and only if $v(p) \in x$ for all $x \in X_H$. In [7], we proved by formula induction the *Key Lemma* stating that for any IntGC-formula A and $x \in X_H$, $x \models A$ if and only if $v(A) \in x$. This enabled us to prove the Kripke-completeness, that is, an IntGC-formula is provable if and only if it is Kripke-valid.

Int2GC-frames. An Int2GC-frame (or a *Kripke frame of Int2GC*) is a quadruple $\mathcal{F} = (X, \leq, R_1, R_2)$ such that X is a non-empty set, $\leq$ is a preorder on X , and R_1 and R_2 are relations on X satisfying

$$(R2) \quad (\geq \circ R_1 \circ \geq) \subseteq R_1$$

$$(R3) \quad (\leq \circ R_2 \circ \leq) \subseteq R_2.$$

Our next lemma is obvious.

Lemma 4.2. *$(X, \leq, R_1, R_2)$ is an Int2GC-frame if and only if $(X, \leq, R_1)$ and $(X, \leq, R_2^{-1})$ are IntGC-frames.*

In Int2GC-frames the valuations and the satisfiability relation $\models$ for $\vee, \wedge, \rightarrow$, and $\neg$ are defined as earlier, but satisfiability of formulas $\blacktriangle A, \nabla A, \triangle A$, and $\blacktriangledown A$ are defined by

$$\begin{aligned} x \models \blacktriangle A &\iff \text{exists } y \text{ such that } x R_1 y \text{ and } y \models A, \\ x \models \nabla A &\iff \text{for all } y, y R_1 x \text{ implies } y \models A, \\ x \models \triangle A &\iff \text{exists } y \text{ such that } y R_2 x \text{ and } y \models A, \text{ and} \\ x \models \blacktriangledown A &\iff \text{for all } y, x R_2 y \text{ implies } y \models A. \end{aligned}$$

It is obvious that Int2GC is *Kripke-sound*, that is, every formula provable in Int2GC is Kripke valid.

We can introduce two IntGC-logics, one with the operators $\blacktriangle$ and ∇ , and the other with $\triangle$ and $\blacktriangledown$. We denote these by IntGC₁ and IntGC₂, respectively. Next we show that Int2GC extends IntGC₁ and IntGC₂.

Lemma 4.3. *Let $\mathcal{F}_i = (X, \leq, R_i)$ be a Kripke-frame for IntGC_i and let A_i be a well-formed formula of IntGC_i, where $i = 1, 2$. Then, A_i is valid in $\mathcal{F}_i$ if and only if A_i is valid in $\mathcal{F} = (X, \leq, R_1, R_2^{-1})$.*

Proof. We prove the claim by formula induction. Concerning IntGC₁-frames and -formulas, the claim is obvious.

Let v be a valuation for the frame $\mathcal{F}_2 = (X, \leq, R_2)$. Thus, $\mathcal{M}_2 = (\mathcal{F}_2, v)$ is an IntGC₂-model and $\mathcal{M} = (\mathcal{F}, v)$ is an Int2GC-model.

Let A_2 be a formula of IntGC_2 which is of the form ΔA for some IntGC_2 -formula A having this property. Then, for all $x \in X$,

$$\begin{aligned} \mathcal{M}_2, x \models \Delta A &\iff (\exists y) x R_2 y \text{ and } \mathcal{M}_2, y \models A \\ &\iff (\exists y) x R_2 y \text{ and } \mathcal{M}, y \models A \\ &\iff (\exists y) y R_2^{-1} x \text{ and } \mathcal{M}, y \models A \\ &\iff \mathcal{M}, x \models \Delta A \end{aligned}$$

Therefore, A_2 is valid in $\mathcal{F}_2$ if and only if A_2 is valid in $\mathcal{F}$. The claim concerning the operator ∇ can be proved analogously. $\square$

The *canonical Int2GC-frame* of an H2GC-algebra $(\mathbb{H}, \triangleright, \triangleleft, \triangleright, \triangleleft)$ is a structure $(X_H, \subseteq, R_1^H, R_2^H)$, where X_H is the set of lattice-filters of H and the relations R_1^H and R_2^H are defined by

$$(x, y) \in R_1^H \iff y \subseteq [x]^{\triangleright^{-1}} \quad \text{and} \quad (x, y) \in R_2^H \iff [x]^{\triangleleft^{-1}} \subseteq y$$

where $[x]^{\triangleleft^{-1}} = \{a \in H \mid a^{\triangleleft} \in x\}$. Equivalently, these relations can be defined as

$$(x, y) \in R_1^H \iff [y]^{\triangleleft^{-1}} \subseteq x \quad \text{and} \quad (x, y) \in R_2^H \iff x \subseteq [y]^{\triangleright^{-1}}$$

in which $[y]^{\triangleright^{-1}} = \{a \in H \mid a^{\triangleright} \in y\}$. The next lemma is obvious and its proof is omitted.

Lemma 4.4. $(\triangleleft^{-1}, \triangleright^{-1})$ and $(\triangleright^{-1}, \triangleleft^{-1})$ are Galois connections on $(\mathcal{O}(H), \subseteq)$.

Similarly, as in the case of HGC-algebras, we can show that the canonical frame $\mathcal{F}^H = (X_H, \subseteq, R_1^H, R_2^H)$ of any H2GC-algebra $(\mathbb{H}, \triangleright, \triangleleft, \triangleright, \triangleleft)$ is an Int2GC -frame. For any valuation v on H , we can define the valuation v^* for the canonical frame $\mathcal{F}^H$ by setting $x \in v^*(p)$ if and only if $v(p) \in x$ for all propositional variables $p \in \text{Var}$ and $x \in X_H$. As in case of IntGC (see Lemma 5.7 in [7]), we can prove by formula induction that the Key Lemma holds, that is, for any Int2GC -formula A and $x \in X_H$, $x \models A$ if and only if $v(A) \in x$. Therefore, we may state the Kripke-completeness presented in the next theorem.

Theorem 4.5 (Completeness II for Int2GC). *A formula A is provable in Int2GC if and only if A is Kripke-valid.*

Proof. Suppose that an Int2GC -formula A is not provable. This means that there exists an H2GC-algebra $(\mathbb{H}, \triangleright, \triangleleft, \triangleright, \triangleleft)$ and a valuation $v: \text{Var} \rightarrow H$ such that $v(A) \neq 1$. We construct the canonical frame $\mathcal{F}^H$ and the valuation v^* as above. Because $v(A) \neq 1$, there exists a prime filter x such that $v(A) \notin x$. By the Key Lemma, this means that $x \not\models A$ in the canonical model $(\mathcal{F}^H, v^*)$. Therefore, A is not Kripke-valid. $\square$

Int2GC+FS-frames. An Int2GC+FS -frame (of a *Kripke frame of Int2GC+FS*) is a triple $\mathcal{F} = (X, \leq, R)$, where X is a non-empty set, $\leq$ is a preorder on X , and R is a relation on X satisfying

$$(R4) \quad (R \circ \leq) \subseteq (\leq \circ R)$$

$$(R5) \quad (\geq \circ R) \subseteq (R \circ \geq).$$

Lemma 4.6. $(X, \leq, R)$ is an Int2GC+FS -frame if and only if $(X, \leq, R \circ \geq, \leq \circ R)$ is an Int2GC -frame.

Proof. Suppose $(X, \leq, R)$ is an Int2GC+FS-frame. Let $R_1 = R \circ \geq$ and $R_2 = \leq \circ R$. We show that the relations R_1 and R_2 satisfy conditions (R2) and (R3). Now $\geq \circ R_1 \circ \geq = \geq \circ (R \circ \geq) \circ \geq = \geq \circ R \circ \geq = (\geq \circ R) \circ \geq \subseteq (R \circ \geq) \circ \geq = R \circ \geq = R_1$, that is, (R2) is satisfied. Similarly, $\leq \circ R_2 \circ \leq = \leq \circ (\leq \circ R) \circ \leq = \leq \circ R \circ \leq = \leq \circ (R \circ \leq) \subseteq \leq \circ (\leq \circ R) = \leq \circ R = R_2$, and also (R3) holds. Thus, $(X, \leq, R \circ \geq, \leq \circ R)$ is an Int2GC-frame.

Conversely, suppose that $(X, \leq, R \circ \geq, \leq \circ R)$ is an Int2GC-frame. We again put $R_1 = R \circ \geq$ and $R_2 = \leq \circ R$. Then, $R \circ \leq \subseteq \leq \circ R \circ \leq = R_2 \circ \leq \subseteq \leq \circ R_2 \circ \leq \subseteq R_2 = \leq \circ R$, that is, (R4) holds. Similarly, $\geq \circ R \subseteq \geq \circ R \circ \geq = \geq \circ R_1 \subseteq \geq \circ R_1 \geq \subseteq R_1 = R \circ \geq$. Hence, also (R5) is satisfied and $(X, \leq, R)$ is an Int2GC+FS-frame. $\square$

Corollary 4.7. *$(X, \leq, R)$ is an Int2GC+FS-frame if and only if $(X, \leq, R \circ \geq)$ and $(X, \leq, R^{-1} \circ \geq)$ are IntGC-frames*

Proof. The claim follows directly from Lemmas 4.2 and 4.6, because $(\leq \circ R)^{-1} = R^{-1} \circ \geq$. $\square$

Again, in Int2GC+FS-frames the valuations and the satisfiability relation $\models$ for $\vee$, $\wedge$, $\rightarrow$, and $\neg$ are defined as earlier, and satisfiability of $\blacktriangle A$, ∇A , $\triangle A$, and $\blacktriangledown A$ are defined by

$$\begin{aligned} x \models \blacktriangle A &\iff \text{exists } y \text{ such that } x(R \circ \geq) y \text{ and } y \models A \\ x \models \nabla A &\iff \text{for all } y, y(R \circ \geq) x \text{ implies } y \models A \\ x \models \triangle A &\iff \text{exists } y \text{ such that } y(\leq \circ R) x \text{ and } y \models A \\ x \models \blacktriangledown A &\iff \text{for all } y, x(\leq \circ R) y \text{ implies } y \models A \end{aligned}$$

Lemma 4.8. *For all Int2GC+FS-models $\mathcal{M} = (\mathcal{F}, v)$ and formulas $A \in \Phi$:*

$$x \models A \text{ and } x \leq y \text{ imply } y \models A.$$

Proof. As an example, we show the claim for $\triangle$ and $\blacktriangledown$.

($\triangle A$) Suppose $x \models \triangle A$ and $x \leq y$. Then, there exists z such that $z(\leq \circ R)x$ and $z \models A$. Thus, there is w such that $z \leq w$, wRx , and $z \models A$. Now wRx and $x \leq y$ imply $w(R \circ \leq)y$. From frame condition (R4), we get $w(\leq \circ R)y$. Now $z \leq w$ implies $z(\leq \circ R)y$. Since $z \models A$, we have $y \models \triangle A$.

($\blacktriangledown A$) Assume that $x \models \blacktriangledown A$, $x \leq y$, but $y \not\models \blacktriangledown A$. Then, there exists z such that $y(\leq \circ R)z$ and $z \not\models A$. Since $x \leq y$, we have $x(\leq \circ R)z$. By $z \not\models A$, we get $x \not\models \blacktriangledown A$, a contradiction. $\square$

Our next lemma showing a connection between validity in Int2GC+FS-frames and Int2GC-frames is obvious and thus its proof is omitted.

Lemma 4.9. *Let $A \in \Phi$. Then, A is valid in the Int2GC+FS-frame $(X, \leq, R)$ if and only A is valid in the Int2GC-frame $(X, \leq, R \circ \geq, \leq \circ R)$.*

Theorem 4.10 (Soundness II for Int2GC+FS). *Every formula provable in Int2GC+FS is Kripke-valid.*

Proof. Suppose that $\triangle A \rightarrow B$ is valid in a Int2GC+FS-frame $(X, \leq, R)$. Then, by Lemma 4.9, $\triangle A \rightarrow B$ is valid in the Int2GC-frame $(X, \leq, R \circ \geq, \leq \circ R)$. This implies that $A \rightarrow \blacktriangledown B$ is valid in the Int2GC-frame $(X, \leq, R \circ \geq, \leq \circ R)$, because Int2GC-preservation of validity of the Galois connection rules. By Lemma 4.9, $A \rightarrow \blacktriangledown B$ is valid in the Int2GC+FS-frame $(X, \leq, R)$. Thus, $(GC \triangle \blacktriangledown)$ preserves validity. Rules $(GC \blacktriangledown \triangle)$, $(GC \blacktriangle \nabla)$, and $(GC \nabla \blacktriangle)$ may be considered similarly.

We show that axiom (D1) is a valid formula. Validity of (D2) can be proved analogously. By Corollary 3.9, this gives that the axioms of Int2GC+FS are valid.

Suppose $x \models \blacktriangle A \wedge \blacktriangledown B$. Then, $x \models \blacktriangle A$ and $x \models \blacktriangledown B$. So, there exists y such that $x(R \circ \geq)y$ and $y \models A$. Thus, there is w such that xRw and $w \geq y$. Because

of persistency, we have $w \models A$. Now $x \leq x$ and xRw imply $x(\leq \circ R)w$. The fact $x \models \nabla B$ means that for all z , $x(\leq \circ R)z$ implies $z \models B$. Therefore, $w \models B$ and thus $w \models A \wedge B$. Because xRw and $w \geq w$, we have $x(R \circ \geq)w$ implying $x \models \blacktriangle(A \wedge B)$. So, (D1) is a valid formula. $\square$

Example 4.11. We present an application showing how preference relations may be used for obtaining particular Kripke-frames of Int2GC+FS . Several definitions of preference structures can be found in the literature; see [14]. There are two fundamental relations, namely “better” (*strict preference*) and “similar” (*indifference*). Here we denote “ b is better than a ” by $a \prec b$ and $a \sim b$ denotes that a and b are similar. Usually, it is assumed that $\prec$ and $\sim$ have at least the following properties:

- (i) $a \prec b$ implies $b \not\prec a$ (*asymmetry of $\prec$*)
- (ii) $a \sim a$ (*reflexivity of $\sim$*)
- (iii) $a \sim b$ implies $b \sim a$ (*symmetry of $\sim$*)
- (iv) $a \prec b$ implies $a \not\sim b$ (*incompatibility of $\prec$ and $\sim$*)

Suppose now that $\prec$ is a transitive strict preference relation on some universe of discourse U . Transitivity is a quite natural property of strict preference, because if a is better than b and b is better than c , also a should be better than c .

Let us denote by $\preceq$ the relation $\prec \cup \Delta_U$, where Δ_U is the *identity relation* of U , that is, $\Delta_U = \{(x, x) \mid x \in U\}$. The relation $\preceq$ is obviously a preorder. Note that since $\prec$ is assumed to be asymmetric, then $a \preceq b$ and $b \preceq a$ imply $a = b$. This means that $\preceq$ is a partial order on U . Assume also that $\preceq$ and $\sim$ are connected by conditions (R4) and (R5), that is,

$$(\sim \circ \preceq) \subseteq (\preceq \circ \sim) \quad \text{and} \quad (\succeq \circ \sim) \subseteq (\sim \circ \succeq),$$

where $\succeq$ is the inverse relation of $\preceq$. These assumptions hold for instance in such object sets which can be organized in “levels” as in Figure 1 – elements in the same level are all similar with respect to their properties, and the elements in an upper level are better than the lower ones.




FIGURE 1.

Hence, the triple $(U, \preceq, \sim)$ can be viewed as an Int2GC+FS -frame. Because the relation $\sim$ is symmetric, ∇A and ∇A have equal interpretations, and the same holds for $\blacktriangle A$ and $\triangle A$. This means that $\nabla A \leftrightarrow \nabla A$ and $\blacktriangle A \leftrightarrow \triangle A$ are valid formulas in any Kripke-model based on the frame $(U, \preceq, \sim)$. This implies, for instance, that $A \rightarrow \nabla \blacktriangle A$ and $\blacktriangle \nabla A \rightarrow A$ are valid in all such Kripke-models for all Int2GC+FS -formulas A .

Additionally, because $\sim$ and $\preceq$ are reflexive, we have that $A \rightarrow \blacktriangle A$ and $\nabla A \rightarrow A$ are valid in all Kripke-models based on $(U, \preceq, \sim)$. Let the formula A represent some property, that is, $x \models A$ means that the object $x \in U$ has this property. The formulas $\blacktriangle A$ and ∇A have the following interpretations:

- (i) $x \models \blacktriangle A$ if there exist $y, z \in U$ such that $x \sim y$, $y \succeq z$ and $z \models A$, that is, x is similar to an object that is better than or equal to an object having the property A .
- (ii) $x \models \blacktriangledown A \iff$ for all y, z , $x \preceq y$ and $y \sim z$ imply $z \models A$, that is, all objects similar to the objects being better or equal to x have the property A .

Thus, the semantics based on preference $\succeq$ and similarity $\sim$ validates many formulas that are not generally **Int2GC+FS**-provable. To get a full correspondence, one should admit R to be any *information relation* satisfying (R4) and (R5), that is,

$$(R \circ \preceq) \subseteq (\preceq \circ R) \quad \text{and} \quad (\succeq \circ R) \subseteq (R \circ \succeq),$$

not just symmetric ones. Various information relations are studied in [4], for instance.

4.2. Canonical Frames and Completeness of Int2GC+FS. To prove the completeness theorem with respect to Kripke-models, we will apply canonical frames and the algebraic completeness for **Int2GC+FS** presented in Theorem 3.8(b).

The *canonical Int2GC+FS-frame* of an **H2GC+FS**-algebra $(\mathbb{H}, \blacktriangleright, \blacktriangleleft, \blacktriangleright, \blacktriangleleft)$ is a structure $(X_H, \subseteq, R^H)$, where X_H is the set of lattice-filters of H and the relation R^H is defined by

$$(x, y) \in R^H \iff [x]^{\blacktriangleleft^{-1}} \subseteq y \subseteq [x]^{\blacktriangleright^{-1}}.$$

The relation R^H can be described also as

$$(x, y) \in R^H \iff [y]^{\blacktriangleleft^{-1}} \subseteq x \subseteq [y]^{\blacktriangleright^{-1}}.$$

This means that $R^H = R_1^H \cap R_2^H$, where R_1^H and R_2^H are the relations of the canonical **Int2GC**-frame of an **H2GC**-algebra.

Next, we will show that the canonical **Int2GC+FS**-frame is a Kripke-frame of **Int2GC+FS**. Before that, we present some results and observations that are needed for our proofs. We denote by $[S]$ the lattice-filter generated by $S \subseteq H$. It is well known that $[S]$ is the set of all elements $a \in H$ such that $a_1 \wedge \dots \wedge a_n \leq a$ for some elements $a_1, \dots, a_n \in S$. We also denote for any $x \in X_H$:

$$\begin{aligned} [x]^{\blacktriangleleft} &= \{a^{\blacktriangleleft} \mid a \in x\}; & [x]^{\blacktriangleright} &= \{a^{\blacktriangleright} \mid a \in x\}; \\ [x]^{\blacktriangleleft} &= \{a^{\blacktriangleleft} \mid a \in x\}; & [x]^{\blacktriangleright} &= \{a^{\blacktriangleright} \mid a \in x\}. \end{aligned}$$

Lemma 4.12. *Let $(\mathbb{H}, \blacktriangleright, \blacktriangleleft, \blacktriangleright, \blacktriangleleft)$ be an **H2GC+FS**-algebra. If k is a filter and y is a prime filter such that $k \cap [-y]^{\blacktriangleleft} = \emptyset$, then there exists a prime filter u such that $k \subseteq u$ and $u \cap [-y]^{\blacktriangleleft} = \emptyset$.*

Proof. Let us denote $\Gamma = \{t \mid t \text{ is a filter, } k \subseteq t, \text{ and } t \cap [-y]^{\blacktriangleleft} = \emptyset\}$. Clearly $\Gamma \neq \emptyset$ and, by Zorn's Lemma, Γ has a maximal element u . Then, u is a filter, $k \subseteq u$, and $u \cap [-y]^{\blacktriangleleft} = \emptyset$.

Assume that u is not a prime filter. Then there exists two elements $a, b \in H$ such that $a \vee b \in u$, but $a, b \notin u$. By maximality of u , this implies that $[u \cup \{a\}]$ and $[u \cup \{b\}]$ are not in Γ . Therefore, we must have that $[u \cup \{a\}] \cap [-y]^{\blacktriangleleft} \neq \emptyset$ and $[u \cup \{b\}] \cap [-y]^{\blacktriangleleft} \neq \emptyset$. So, there exists $c, d \in [-y]^{\blacktriangleleft}$ such that $c \in [u \cup \{a\}]$ and $d \in [u \cup \{b\}]$. Because u is a filter, this implies that there exist $e, f \in u$ such that $e \wedge a \leq c$ and $f \wedge b \leq d$. Now $e \wedge f \in u$ and $a \vee b \in u$ imply $(e \wedge f) \wedge (a \vee b) \in u$. Since

$$(e \wedge f) \wedge (a \vee b) = (e \wedge f \wedge a) \vee (e \wedge f \wedge b) \leq c \vee d,$$

we obtain $c \vee d \in u$. Now there exist $c_1, d_1 \in -y$ such that $c = c_1^{\blacktriangleleft}$ and $d = d_1^{\blacktriangleleft}$. Because $c \vee d \in u$ and $c \vee d = c_1^{\blacktriangleleft} \vee d_1^{\blacktriangleleft} \leq (c_1 \vee d_1)^{\blacktriangleleft}$, we have $(c_1 \vee d_1)^{\blacktriangleleft} \in u$. On the other hand, $c_1, d_1 \notin y$ implies $c_1 \vee d_1 \notin y$, because y is a prime filter. Thus, $c_1 \vee d_1 \in -y$ implies $(c_1 \vee d_1)^{\blacktriangleleft} \in [-y]^{\blacktriangleleft}$. But $u \cap [-y]^{\blacktriangleleft} = \emptyset$, a contradiction. Thus, u is a prime filter. $\square$

Let S be a non-empty subset of a lattice L such that $a \vee b \in S$ implies $a \in S$ or $b \in S$ for all $a, b \in L$. It is easily seen that such sets S can be characterised as the sets whose set-theoretical complement $-S$ is a $\vee$ -subsemilattice of L . In [7], we proved the following lemma.

Lemma 4.13. *Let L be a distributive lattice. If x is a filter and u is a superset of x such that its set-theoretical complement $-u$ is a $\vee$ -subsemilattice of L , then there exists a prime filter z such that $x \subseteq z \subseteq u$.*

Our next proposition shows that the canonical frames are $\text{Int2GC}+\text{FS}$ -frames.

Proposition 4.14. *If $(\mathbb{H}, \triangleright, \triangleleft, \triangleright, \triangleleft)$ is an $H2GC+\text{FS}$ -algebra, then $(X^H, \subseteq, R^H)$ is an $\text{Int2GC}+\text{FS}$ -frame.*

Proof. (R4) Assume that $x(R^H \circ \subseteq)y$. This implies that there exists $z \in X_H$ such that $x R^H z$ and $z \subseteq y$. Then, $[x]^{\triangleleft^{-1}} \subseteq z \subseteq [x]^{\triangleright^{-1}}$. Let $k = [x \cup [y]^{\triangleright}]$ be the filter generated by $x \cup [y]^{\triangleright}$. We show first that $k \cap [-y]^{\triangleleft} = \emptyset$. Namely, if $k \cap [-y]^{\triangleleft} \neq \emptyset$, then there exists an element a such that $a \in k$ and $a \in [-y]^{\triangleleft}$. Since $a \in k = [x \cup [y]^{\triangleright}]$, there are $b \in x$ (recall that x is a filter) and $c_1, \dots, c_n \in y$ such that $b \wedge c_1^{\triangleright} \wedge \dots \wedge c_n^{\triangleright} \leq a$. Let us denote $c = c_1 \wedge \dots \wedge c_n \in y$. Because the map $\triangleright$ is order-preserving, we have $c^{\triangleright} \leq c_1^{\triangleright} \wedge \dots \wedge c_n^{\triangleright}$. This means that $b \wedge c^{\triangleright} \leq a$ and $b \leq c^{\triangleright} \rightarrow a$. Now $a = d^{\triangleleft}$ for some $d \in -y$, and so $b \leq c^{\triangleright} \rightarrow d^{\triangleleft} \leq (c \rightarrow d)^{\triangleleft}$ by (fs3). Since $b \in x$ and x is a filter, we have $(c \rightarrow d)^{\triangleleft} \in x$. From this we get $c \rightarrow d \in [x]^{\triangleleft^{-1}} \subseteq z \subseteq y$. Because $c \in y$ and y is a filter, also $c \wedge (c \rightarrow d) \in y$. Since $c \wedge (c \rightarrow d) \leq d$, we have $d \in y$, a contradiction. Hence, $k \cap [-y]^{\triangleleft} = \emptyset$.

By Lemma 4.12, there exists a prime filter u such that $k \subseteq u$ and $u \cap [-y]^{\triangleleft} = \emptyset$. So, $x \subseteq k \subseteq u$, $[y]^{\triangleright} \subseteq k \subseteq u$, and $y \subseteq [u]^{\triangleright^{-1}}$. In addition, $[u]^{\triangleleft^{-1}} \subseteq y$, because if $a \in [u]^{\triangleleft^{-1}}$, then $a^{\triangleleft} \in u$ which gives $a^{\triangleleft} \notin [-y]^{\triangleleft}$, because $u \cap [-y]^{\triangleleft} = \emptyset$. Thus, $a \notin -y$, that is, $a \in y$. Now $x \subseteq u$ and $u R^H y$ give $x(\subseteq \circ R^H)y$.

(R5) Assume $x(\supseteq \circ R^H)y$. Then for some $w \in X_H$, $x \supseteq w$ and $[w]^{\triangleleft^{-1}} \subseteq y \subseteq [w]^{\triangleright^{-1}}$. Hence, $y \subseteq [x]^{\triangleright^{-1}}$, because $\triangleright^{-1}$ is order-preserving. To show that $x(R^H \circ \supseteq)y$, we need to find a prime filter $z \in X_H$ such that $[x]^{\triangleleft^{-1}} \subseteq z \subseteq [x]^{\triangleright^{-1}}$ and $z \supseteq y$. Consider the filter $k = [y \cup [x]^{\triangleleft^{-1}}]$. We show first that $k \subseteq [x]^{\triangleright^{-1}}$. Assume $a \in k$. Then, there exists $c \in y$ and $d \in [x]^{\triangleleft^{-1}}$ such that $c \wedge d \leq a$ (note that y is a filter and $[x]^{\triangleleft^{-1}}$ is closed under meets). Hence, $c \leq d \rightarrow a$ and $c^{\triangleright} \leq (d \rightarrow a)^{\triangleright} \leq d^{\triangleleft} \rightarrow a^{\triangleright}$ by (fs1). Since $c \in y \subseteq [x]^{\triangleright^{-1}}$, we have $c^{\triangleright} \in x$ and $d^{\triangleleft} \rightarrow a^{\triangleright} \in x$. Because $d^{\triangleleft} \in x$, we obtain $a^{\triangleright} \in x$, that is, $a \in [x]^{\triangleright^{-1}}$ as required.

Because $k \subseteq [x]^{\triangleright^{-1}}$, k is a filter, and $[x]^{\triangleright^{-1}}$ is a set such that its set-theoretical complement is a $\vee$ -subsemilattice of L , by Lemma 4.13 there exists $z \in X_H$ such that $k \subseteq z \subseteq [x]^{\triangleright^{-1}}$. Combining the above observations, we have $z \supseteq k \supseteq y$ and $[x]^{\triangleleft^{-1}} \subseteq k \subseteq z \subseteq [x]^{\triangleright^{-1}}$, that is, $x R^H z$ and $z \supseteq y$. $\square$

Let $(\mathbb{H}, \triangleright, \triangleleft, \triangleright, \triangleleft)$ be an $H2GC+\text{FS}$ -algebra. Let $v: \text{Var} \rightarrow H$ be a valuation. We may now define a valuation $v^*: \text{Var} \rightarrow X_H$ for the canonical frame $\mathcal{F}^H = (X_H, \subseteq, R^H)$ by setting $x \in v^*(p)$ if and only if $v(p) \in x$ for all $p \in \text{Var}$. Hence, in the canonical model $(\mathcal{F}^H, v^*)$, we have $x \models p$ if and only if $v(p) \in x$ for all $x \in X_H$. We show that an analogous condition holds for all formulas A .

Lemma 4.15 (Key Lemma). *Let $(\mathbb{H}, \triangleright, \triangleleft, \triangleright, \triangleleft)$ be an $H2GC+\text{FS}$ -algebra and $v: \text{Var} \rightarrow H$ a valuation. In the canonical model $(\mathcal{F}^H, v^*)$, we have $x \models A$ if and only if $v(A) \in x$ for all $x \in X_H$ and $A \in \Phi$.*

Proof. We prove the result by formula induction. For the connectives $\vee, \wedge, \rightarrow$, and $\neg$ the result is well known from the theory of intuitionistic logic. In addition, we

only show the proofs for formulas ΔA and ∇A , since for $\blacktriangle A$ and ∇A the proofs are analogous.

($\blacktriangle A$) Suppose that $x \models \blacktriangle A$. This means that there exists a prime filter y such that $x (R^H \circ \supseteq) y$ and $y \models A$. By the induction hypothesis, we have that $y \in v(A)$. In addition, there exists a prime filter u such that $x R^H u$ and $y \subseteq u$, that is, $[x]^{\blacktriangleleft^{-1}} \subseteq u \subseteq [x]^{\blacktriangleright^{-1}}$. We obtain directly that $v(A) \in y \subseteq u \subseteq [x]^{\blacktriangleright^{-1}}$, which means that $v(A)^{\blacktriangleright} = v(\blacktriangle A) \in x$.

Conversely, suppose $v(\blacktriangle A) = v(A)^{\blacktriangleright} \in x$. Let us consider the filter $k = [x]^{\blacktriangleleft^{-1}} \cup \{v(A)\}$. First we show that $k \subseteq [x]^{\blacktriangleright^{-1}}$. Assume that $a \in k$. Then there exists $b \in [x]^{\blacktriangleleft^{-1}}$ such that $b \wedge v(A) \leq a$ (note that $[x]^{\blacktriangleleft^{-1}}$ is closed under finite meets). We have that $v(A) \leq b \rightarrow a$ and $v(A)^{\blacktriangleright} \leq (b \rightarrow a)^{\blacktriangleright} \leq b^{\blacktriangleleft} \rightarrow a^{\blacktriangleright}$ by (fs1). This implies $b^{\blacktriangleleft} \rightarrow a^{\blacktriangleright} \in x$. Because $b^{\blacktriangleleft} \in x$, we obtain $a^{\blacktriangleright} \in x$ and $a \in [x]^{\blacktriangleright^{-1}}$. Hence, $k \subseteq [x]^{\blacktriangleright^{-1}}$. Because k is a filter and $[x]^{\blacktriangleright^{-1}}$ is a set such that its set-theoretical complement is a $\vee$ -subsemilattice of H , we have by Lemma 4.13 that there exists a prime filter y such that $k \subseteq y \subseteq [x]^{\blacktriangleright^{-1}}$. By the definition of k , $[x]^{\blacktriangleleft^{-1}} \subseteq k \subseteq y$ and $v(A) \in k \subseteq y$. We have $[x]^{\blacktriangleleft^{-1}} \subseteq y \subseteq [x]^{\blacktriangleright^{-1}}$, that is, $x R^H y$. By the induction hypothesis, $y \models A$. Since $y \supseteq y$ holds trivially, we have $x (R^H \circ \supseteq) y$ implying $x \models \blacktriangle A$.

(∇A) Suppose that $v(\nabla A) = v(A)^{\blacktriangleleft} \in x$. Let $y \in X_H$. If $y (R^H \circ \supseteq) x$, then there exists z such that $y R^H z$ and $z \supseteq x$. Now $y R^H z$ is equivalent to $[z]^{\blacktriangleleft^{-1}} \subseteq y \subseteq [z]^{\blacktriangleright^{-1}}$. Therefore, $v(A)^{\blacktriangleleft} \in x \subseteq z$ gives $v(A) \in [z]^{\blacktriangleleft^{-1}} \subseteq y$. By the induction hypothesis, $y \models A$ and hence $x \models \nabla A$.

For the other direction, assume $v(\nabla A) = v(A)^{\blacktriangleleft} \notin x$, that is, $v(A) \notin [x]^{\blacktriangleleft^{-1}}$. It is easy to observe that $[x]^{\blacktriangleleft^{-1}}$ is a filter. Then, by the Prime Filter Theorem of distributive lattices (see [7, Lemma 5.4], for instance), there exists a prime filter u such that $v(A) \notin u$ and $[x]^{\blacktriangleleft^{-1}} \subseteq u$.

Let us consider the filter $k = [x \cup [u]^{\blacktriangleright}]$. We first show that $k \cap [-u]^{\blacktriangleleft} = \emptyset$. If $k \cap [-u]^{\blacktriangleleft} \neq \emptyset$, then there exists $a \in k \cap [-u]^{\blacktriangleleft}$. Because $a \in k = [x \cup [u]^{\blacktriangleright}]$, there are $b \in x$ and $c_1, \dots, c_n \in u$ such that $b \wedge c_1^{\blacktriangleright} \wedge \dots \wedge c_n^{\blacktriangleright} \leq a$ (recall that x is a filter). Let us denote $c = c_1 \wedge \dots \wedge c_n \in u$. Hence, $c^{\blacktriangleright} \leq c_1^{\blacktriangleright} \wedge \dots \wedge c_n^{\blacktriangleright}$ and $b \wedge c^{\blacktriangleright} \leq a$. But now $a = d^{\blacktriangleleft}$ for some $d \notin u$. So, $b \wedge c^{\blacktriangleright} \leq d^{\blacktriangleleft}$. This gives that $b \leq c^{\blacktriangleright} \rightarrow d^{\blacktriangleleft} \leq (c \rightarrow d)^{\blacktriangleleft}$ by (fs4). Because $b \in x$, we have $(c \rightarrow d)^{\blacktriangleleft} \in x$. This means that $c \rightarrow d \in [x]^{\blacktriangleleft^{-1}} \subseteq u$. Now $c \in u$ implies $d \in u$, a contraction. Therefore, $k \cap [-u]^{\blacktriangleleft} = \emptyset$.

By Lemma 4.12, there exists a prime filter y such that $k \subseteq y$ and $y \cap [-u]^{\blacktriangleleft} = \emptyset$. Since $k \subseteq y$, we have $x \subseteq y$ and $[u]^{\blacktriangleright} \subseteq y$ meaning $u \subseteq [y]^{\blacktriangleright^{-1}}$. The fact that $y \cap [-u]^{\blacktriangleleft} = \emptyset$ implies $[y]^{\blacktriangleleft^{-1}} \subseteq u$, because if $a \in [y]^{\blacktriangleleft^{-1}}$, then $a^{\blacktriangleleft} \in y$. This gives $a^{\blacktriangleleft} \notin [-u]^{\blacktriangleleft}$, $a \notin -u$, and $a \in u$. By combining our observations, we have $[y]^{\blacktriangleleft^{-1}} \subseteq u \subseteq [y]^{\blacktriangleright^{-1}}$, that is, $u R^H y$ and $y \supseteq x$. Thus, $u (R^H \circ \supseteq) x$. Because $v(A) \notin u$, we have $u \not\models A$ by the induction hypothesis. Hence, $x \not\models \nabla A$. $\square$

As in case of Theorem 4.5, we may prove the following completeness result by applying the Key Lemma.

Theorem 4.16 (Completeness II for Int2GC+FS). *A formula A is provable in Int2GC+FS if and only if A is Kripke-valid.*

5. CONNECTIONS TO INTUITIONISTIC TENSE LOGIC

Intuitionistic tense logic $\mathbf{IK}_t$ was introduced by Ewald [9] by extending the language of intuitionistic propositional logic with the usual temporal expressions FA (A is true at some future time), PA (A was true at some past time), GA (A will

be true at all future times), and HA (A has always been true in the past). The Hilbert-style axiomatisation of IK_t can be found in [9, p. 171]:

- | | |
|---|--|
| (1) All axioms of intuitionistic logic | |
| (2) $G(A \rightarrow B) \rightarrow (GA \rightarrow GB)$ | (2') $H(A \rightarrow B) \rightarrow (HA \rightarrow HB)$ |
| (3) $G(A \wedge B) \leftrightarrow GA \wedge GB$ | (3') $H(A \wedge B) \leftrightarrow HA \wedge HB$ |
| (4) $F(A \vee B) \leftrightarrow FA \vee FB$ | (4') $P(A \vee B) \leftrightarrow PA \vee PB$ |
| (5) $G(A \rightarrow B) \rightarrow (FA \rightarrow FB)$ | (5') $H(A \rightarrow B) \rightarrow (PA \rightarrow PB)$ |
| (6) $GA \wedge FB \rightarrow F(A \wedge B)$ | (6') $HA \wedge PB \rightarrow P(A \wedge B)$ |
| (7) $G\neg A \rightarrow \neg FA$ | (7') $H\neg A \rightarrow \neg PA$ |
| (8) $FHA \rightarrow A$ | (8') $PGA \rightarrow A$ |
| (9) $A \rightarrow HFA$ | (9') $A \rightarrow GPA$ |
| (10) $(FA \rightarrow GB) \rightarrow G(A \rightarrow B)$ | (10') $(PA \rightarrow HB) \rightarrow H(A \rightarrow B)$ |
| (11) $F(A \rightarrow B) \rightarrow (GA \rightarrow FB)$ | (11') $P(A \rightarrow B) \rightarrow (HA \rightarrow PB)$ |

The rules of inference are modus ponens (MP), and

$$(\text{RH}) \quad \frac{A}{HA} \qquad (\text{RG}) \quad \frac{A}{GA}$$

Our next proposition shows that if we identify $\blacktriangle, \blacktriangledown, \triangle, \triangledown$ with F, G, P, H , respectively, then $\text{Int2GC}+\text{FS}$ and IK_t will become syntactically equivalent. Recall that

$$\text{Int2GC}+\text{FS} = \text{Int2GC} + \{(\text{FS1}) \text{ or } (\text{FS4}) \text{ or } (\text{D1})\} + \{(\text{FS2}) \text{ or } (\text{FS3}) \text{ or } (\text{D2})\},$$

and Int2GC is obtained by extending intuitionistic logic with rules $(\text{GC} \nabla \blacktriangle)$, $(\text{GC} \blacktriangle \nabla)$, $(\text{GC} \blacktriangledown \triangle)$, and $(\text{GC} \triangle \blacktriangledown)$.

Theorem 5.1. $\mathsf{IK}_t = \text{Int2GC}+\text{FS}$.

Proof. First we show that all axioms IK_t are provable in $\text{Int2GC}+\text{FS}$, and all rules of IK_t are admissible in $\text{Int2GC}+\text{FS}$. In this first part, let $\vdash$ denote that a formula A is provable in $\text{Int2GC}+\text{FS}$. As noted in Section 2, axioms (2), (2'), (3), (3') (4), (4'), (8), (8'), (9), (9') are provable even in Int2GC . Additionally, rules (MP), (RH), and (RG) are admissible in Int2GC . Axioms (10), (10'), (11), (11') are the Fischer Servi axioms (FS3), (FS4), (FS1), (FS2), so they are provable in $\text{Int2GC}+\text{FS}$.

Axiom (FS1) is equivalent to $\blacktriangle(A \rightarrow B) \wedge \nabla A \rightarrow \blacktriangle B$. If we set $B := A \wedge B$ in this formula, we have that $\vdash (\blacktriangle(A \rightarrow A \wedge B) \wedge \nabla A) \rightarrow \blacktriangle(A \wedge B)$. Because $A \rightarrow A \wedge B$ is equivalent to $A \rightarrow B$, and $\vdash B \rightarrow (A \rightarrow B)$ gives $\vdash \blacktriangle B \rightarrow \blacktriangle(A \rightarrow B)$ by the monotonicity of $\blacktriangle$, we obtain $\vdash \blacktriangle B \wedge \nabla A \rightarrow \blacktriangle(A \wedge B)$ and thus (6) is provable in $\text{Int2GC}+\text{FS}$. Provability of (6') can be shown similarly.

Because $\vdash \blacktriangle \nabla(A \rightarrow B) \rightarrow (A \rightarrow B)$, we have $\vdash \blacktriangle \nabla(A \rightarrow B) \wedge A \rightarrow B$ and $\vdash \triangle(\blacktriangle \nabla(A \rightarrow B) \wedge A) \rightarrow \triangle B$. Let us set $A := \blacktriangle \nabla(A \rightarrow B)$ and $B := A$ in axiom (6') (which we just showed to be provable in $\text{Int2GC}+\text{FS}$). We obtain $\vdash \nabla \blacktriangle \nabla(A \rightarrow B) \wedge \triangle A \rightarrow \triangle(\blacktriangle \nabla(A \rightarrow B) \wedge A)$. Thus, $\vdash \nabla \blacktriangle \nabla(A \rightarrow B) \wedge \triangle A \rightarrow \triangle B$. Because $\vdash \nabla(A \rightarrow B) \rightarrow \nabla \blacktriangle \nabla(A \rightarrow B)$, we have $\vdash \nabla(A \rightarrow B) \wedge \triangle A \rightarrow \triangle B$. This is equivalent to $\vdash \nabla(A \rightarrow B) \rightarrow (\triangle A \rightarrow \triangle B)$. Hence, (5') is provable in $\text{Int2GC}+\text{FS}$, and provability of (5) can be showed in an analogous manner.

If we set $B := \perp$ in (5), we get $\vdash \blacktriangledown(A \rightarrow \perp) \rightarrow (\blacktriangle A \rightarrow \blacktriangle \perp)$. Because $\blacktriangle \perp$ is equivalent to $\perp$, we have $\vdash \blacktriangledown \neg A \rightarrow \neg \blacktriangle A$. This means that (7) and (7') are provable.

Because axioms (10), (10'), (11), (11') are the Fischer Servi axioms, for the other direction is enough to show admissibility of rules $(\text{GC} \nabla \blacktriangle)$, $(\text{GC} \blacktriangle \nabla)$, $(\text{GC} \blacktriangledown \triangle)$, $(\text{GC} \triangle \blacktriangledown)$ in IK_t . First, we show admissibility of the rules of monotonicity, that is, if $A \rightarrow B$ is provable, then $HA \rightarrow HB$, $PA \rightarrow PB$, $GA \rightarrow GB$, and $FA \rightarrow FB$ are provable.

Here $\vdash A$ denotes that the formula A is provable in IK_t . Assume $\vdash A \rightarrow B$. By (RG), $\vdash G(A \rightarrow B)$. Now $\vdash GA \rightarrow GB$ follows by (2), and from $\vdash G(A \rightarrow B)$, we obtain also $\vdash FA \rightarrow FB$ by (5). Similarly, $\vdash A \rightarrow B$ implies $\vdash HA \rightarrow HB$ and $\vdash PA \rightarrow PB$ by applying (RH), (2'), and (5').

Next we prove admissibility of $(\mathsf{GC} \nabla \blacktriangle)$. Assume that $\vdash A \rightarrow HB$. Then, $FA \rightarrow FHB$ by monotonicity of F . Because $\vdash FHB \rightarrow B$ by (8), we obtain $\vdash FA \rightarrow B$. Similarly, by (8') and monotonicity of P , $A \rightarrow GB$ implies $PA \rightarrow B$, that is, $(\mathsf{GC} \blacktriangledown \Delta)$ is admissible in IK_t . Monotonicity of H and axiom (9) yield $FA \rightarrow B$ implies $A \rightarrow HB$, and monotonicity of G and (9') give that $PA \rightarrow B$ implies $A \rightarrow BG$. Thus, rules $(\mathsf{GC} \blacktriangle \nabla)$ and $(\mathsf{GC} \Delta \blacktriangledown)$ are admissible. $\square$

Remark 5.2. It is proved in [15] that ILGC is equivalent, with respect to provability, to the minimal (classical) tense logic K_t , that is, ILGC can be viewed as a simple formulation of K_t . The same analogy applies here, because $\mathsf{Int2GC+FS}$ can be seen as an alternative formulation of IK_t .

It should be noted that with respect to Kripke-semantics, IK_t and $\mathsf{Int2GC+FS}$ are quite different. A Kripke-frame of IK_t consists of a partially-ordered set $(\Gamma, \leq)$ (the “states-of-knowledge”), family of sets T_γ , where $\gamma \in \Gamma$ (times known at state-of-knowledge γ), such that $\gamma \leq \varphi$ implies $T_\gamma \subseteq T_\varphi$, meaning that advancing in knowledge retains what is known about times and their temporal ordering, and a collection of binary relations μ_γ on T_γ (the temporal ordering of T_γ as it is understood at state-of-knowledge γ) [9], whereas $\mathsf{Int2GC+FS}$ is conceived as an *information logic* such that its frames $(X, \leq, R)$ are such that X forms the *universe of discourse*, and $\leq$ and R are relations reflecting relationships between the objects in X , such as preference and indifference of objects (see Example 4.11).

It is also obvious and well-known that the axiomatisation of Ewald is not minimal, because several axioms can be deduced from the other axioms. We present a reduced axiomatisation, in which the number of axioms is the half of the size of the axiomatisation in [9].

Proposition 5.3. IK_t can be axiomatised by adding (2), (2'), (5), (5'), (8), (8'), (9), (9'), (11), (11') to the axioms of intuitionistic logic together with rules (MP), (RH), and (RG).

Proof. As shown in the proof of Theorem 5.1, if $\blacktriangle, \blacktriangledown, \Delta, \nabla$ are identified with F, G, P, H , then axioms (2), (2'), (5), (5'), (8), (8'), (9), (9') with rules (MP), (RH) and (RG) are enough to show that rules $(\mathsf{GC} \nabla \blacktriangle), (\mathsf{GC} \blacktriangle \nabla), (\mathsf{GC} \blacktriangledown \Delta), (\mathsf{GC} \Delta \blacktriangledown)$ are admissible. Axioms (11), (11') coincide with (FS1) and (FS2), so the proof is complete, because $\mathsf{IK}_t = \mathsf{Int2GC+FS}$. $\square$

In the next proposition, we present another axiomatisation of $\mathsf{Int2GC}$ using axioms of intuitionistic tense logic.

Proposition 5.4. $\mathsf{Int2GC}$ can be axiomatised by adding (2), (2'), (8), (8'), (9), (9') to the axioms of intuitionistic logic together with rules (MP), (RH), (RG), and rules:

$$\begin{array}{ll} (\mathsf{RMF}) \quad \frac{A \rightarrow B}{FA \rightarrow FB} & (\mathsf{RMP}) \quad \frac{A \rightarrow B}{PA \rightarrow PB} \end{array}$$

Proof. Monotonicity of G and H follow from (RG), (RH), (2), and (2'). Because all operators are thus monotone, admissibility of rules $(\mathsf{GC} \nabla \blacktriangle), (\mathsf{GC} \blacktriangle \nabla), (\mathsf{GC} \blacktriangledown \Delta), (\mathsf{GC} \Delta \blacktriangledown)$ follow easily from (8), (8'), (9), (9').

On the other hand, in Section 2 we have noted that axioms (2), (2'), (8), (8'), (9), (9') are provable in $\mathsf{Int2GC}$ and rules (RH), (RG), (RMF), (RMP) are admissible. $\square$

Let us observe that Int2GC cannot be axiomatised by using only axioms and rules of Ewald's system. The reason for this is that monotonicity of operators P and F need to be added, since rules (RMF) and (RMP) do not belong to the system by Ewald as "initial rules", even they are admissible in $\mathbf{IK}_t$. On the other hand, monotonicity of P and F could be obtained by adding axioms (5) and (5') to the system of Proposition 5.4 (without monotonicity of F and P), but then we have a logic which is too strong, since (5) and (5') cannot be proved in Int2GC – this is because Galois connections $(\blacktriangle, \nabla)$ and $(\Delta, \blacktriangledown)$ are "independent", that is, operations $\blacktriangle$ and $\blacktriangledown$ are not in anyway connected with each other. For instance, consider an H2GC-algebra on the three element chain $0 < u < 1$ such that $\blacktriangleright$ and $\blacktriangleleft$ equal the identity mapping, and $x^\blacktriangleright = 0$ and $x^\blacktriangleleft = 1$ for all $x \in \{0, u, 1\}$. Then $(1 \rightarrow u)^\blacktriangleleft = 1$, but $1^\blacktriangleright \rightarrow u^\blacktriangleright = 1 \rightarrow u = u$. This actually means that we have an "intermediate logic" Int2GC + $\{(5), (5')\}$ situated between Int2GC and Int2GC+FS. However, the study of Int2GC + $\{(5), (5')\}$ is confined outside of the scope of this work.

REFERENCES

- [1] R. Balbes and Ph. Dwinger, *Distributive lattices*, University of Missouri Press, Columbia, Missouri, 1974.
- [2] R. Bělohlávek, *Fuzzy Galois connections*, Mathematical Logic Quarterly **45** (1999), 497–504.
- [3] D. van Dalen, *Intuitionistic logic*, The Blackwell guide to philosophical logic, 2001, pp. 224–257.
- [4] S. P. Demri and E. S. Orłowska, *Incomplete information: Structure, inference, complexity*, Springer-Verlag, Berlin, Heidelberg, 2002.
- [5] D. Dubois and H. Prade, *Rough fuzzy sets and fuzzy rough sets*, International Journal of General Systems **17** (1990), 191–209.
- [6] J. M. Dunn, *Positive modal logic*, Studia Logica **55** (1995), 301–317.
- [7] W. Dzik, J. Järvinen, and M. Kondo, *Intuitionistic propositional logic with Galois connections*, Logic Journal of the IGPL **18** (2010), 837–858.
- [8] M. Ern , J. Koslowski, A. Melton, and G. E. Strecker, *A primer on Galois connections*, Annals of the New York Academy of Sciences **704** (1993), 103–125.
- [9] W. B. Ewald, *Intuitionistic tense and modal logic*, The Journal of Symbolic Logic **51** (1986), 166–179.
- [10] G. Fischer Servi, *Axiomatizations for some intuitionistic modal logics*, Rendiconti del Seminario Matematico dell' Universit  Politecnica di Torino **42** (1984), 179–194.
- [11] G. Georgescu and A. Popescu, *Non-dual fuzzy connections*, Archive for Mathematical Logic **43** (2004), 1009–1039.
- [12] J. A. Goguen, *L-fuzzy sets*, Journal of Mathematical Analysis and Applications **18** (1967), 145–174.
- [13] G. Gr tzer, *General lattice theory*, 2nd ed., Birkh user, Basel, 1998.
- [14] S. O. Hansson and T. Gr ne-Yanoff, *Preferences*, The Stanford encyclopedia of philosophy, 2009. <http://plato.stanford.edu/archives/spr2009/entries/preferences/>.
- [15] J. J rvinen, M. Kondo, and J. Kortelainen, *Logics from Galois connections*, International Journal of Approximate Reasoning **49** (2008), 595–606.
- [16] Z. Pawlak, *Rough sets*, International Journal of Computer and Information Sciences **11** (1982), 341–356.
- [17] H. Rasiowa and R. Sikorski, *The mathematics of metamathematics*, 2nd ed., PWN-Polish Scientific Publishers, Warsaw, 1968.

WOJCIECH DZIK, INSTITUTE OF MATHEMATICS, UNIVERSITY OF SILESIA, UL. BANKOWA 12, 40-007 KATOWICE, POLAND

E-mail address: dzikw@silesia.top.pl

JOUNI J RVINEN, TURKU, FINLAND

E-mail address: Jouni.Kalervo.Jarvinen@gmail.com

MICHIRO KONDO, SCHOOL OF INFORMATION ENVIRONMENT, TOKYO DENKI UNIVERSITY, INZAI, 270-1382, JAPAN

E-mail address: kondo@sie.dendai.ac.jp